

FORM PTO-1390
(REV 10-2000)

U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE

ATTORNEY'S DOCKET NUMBER

R11-003CPUS

**TRANSMITTAL LETTER TO THE UNITED STATES
DESIGNATED/ELECTED OFFICE (DO/EO/US)
CONCERNING A FILING UNDER 35 U.S.C.371**

U.S. APPLICATION NO. (If known, see 37 CFR 1.5)

09/719277

INTERNATIONAL APPLICATION
PCT/US99/12929

INTERNATIONAL FILING DATE
09 June 1999 (09.06.99)

PRIORITY DATE CLAIMED
**09 June 1998 (09.06.98)
11 June 1998 (11.06.98)**

TITLE OF INVENTION

NOVEL HEPATITIS C VIRUS PEPTIDES AND USES THEREOF

APPLICANT(S) FOR DO/EO/US

Andrea B. BRANCH, Jose L. WALEWSKI, and Dechard D. STUMP

Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:

1. ☒ This is a **FIRST** submission of items concerning a filing under 35 U.S.C.371.
2. ☐ This is a **SECOND** or **SUBSEQUENT** submission of items concerning a filing under 35 U.S.C. 371.
3. ☐ This is an express request to promptly begin national examination procedures (35 U.S.C. 371(f)).
4. ☒ The US has been elected by the expiration of 19 months from the priority date (PCT Article 31).
5. ☒ A copy of the International Application as filed (35 U.S.C. 371(c)(2))
 - a. ☐ is attached hereto (required only if not communicated by the International Bureau).
 - b. ☒ has been communicated by the International Bureau.
 - c. ☐ is not required, as the application was filed in the United States Receiving Office (RO/US).
6. ☐ An English language translation of the International Application as filed (35 U.S.C 371(c)(2)).
7. ☒ Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371(c)(3))
 - a. ☐ are attached hereto (required only if not communicated by the International Bureau).
 - b. ☐ have been communicated by the International Bureau.
 - c. ☐ have not been made; however, the time limit for making such amendments has NOT expired.
 - d. ☒ have not been made and will not be made.
8. ☐ An English language translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)).
9. ☒ An oath or declaration of the inventor(s) (35 U.S.C. 371(c)(4)). **(unexecuted) (4 sheets)**
10. ☐ An English language translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)).

Items 11. to 16. below concern document(s) or information included:

11. ☐ An Information Disclosure Statement under 37 CFR 1.97 and 1.98.
12. ☐ An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included
13. ☐ A **FIRST** preliminary amendment.
☐ A **SECOND** or **SUBSEQUENT** preliminary amendment.
14. ☐ A substitute specification.
15. ☐ A change of power of attorney and/or address letter.
16. ☒ Other items or information: **Transmittal Letter (2 sheets in duplicate); PCT International Published Application (WO 99/63941) (without International Search Report) (51 sheets); PCT International Published Application Cover Sheet (WO 99/63941)(with International Search Report attached) (3 sheets); Sequence Listing (3 sheets); Diskette Containing Sequence Listing; Check (\$520) based on small entity status; Certificate of Express Mailing (1 sheet); and Return Postcard.**

U.S. APPLICATION NO. (if known, see 37 CFR 1.5) <div style="font-size: 24pt; font-weight: bold; margin-top: 5px;">09/719277</div>		INTERNATIONAL APPLICATION NO. <div style="font-weight: bold; margin-top: 5px;">PCT/US99/12929</div>		ATTORNEY'S DOCKET NO. <div style="font-weight: bold; margin-top: 5px;">RII-003CPUS</div>	
17. <input checked="" type="checkbox"/> The following fees are submitted:				CALCULATIONS PTO USE ONLY	
BASIC NATIONAL FEE (37 CFR 1.492 (a) (1) - (5)) .(a/o November 1, 2000): Neither international preliminary examination fee (37 CFR 1.482) nor international search fee (37 CFR 1.445(a)(2)) paid to USPTO and International Search Report not prepared by the EPO or JPO.....\$1000 International preliminary examination fee (37 CFR 1.482) not paid to USPTO but International Search Report prepared by the EPO or JPO\$860 International preliminary examination fee (37 CFR 1.482) not paid to USPTO but international search fee (37 CFR 1.455(a)(2)) paid to USPTO\$710 International preliminary examination fee paid to USPTO (37 CFR 1.482) but all claims did not satisfy provisions of PCT Article 33(1)-(4).....\$690 International preliminary examination fee paid to USPTO (37 CFR 1.482) and all claims satisfied provisions of PCT Article 33(1)-(4).....\$100 <div style="text-align: right;">ENTER APPROPRIATE BASIC FEE AMOUNT =</div>				<div style="border: 1px solid black; padding: 5px; display: inline-block;">\$860</div>	
Surcharge of \$130.00 for furnishing the oath or declaration later than <input type="checkbox"/> 20 <input type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492(e)).				<div style="border: 1px solid black; padding: 5px; display: inline-block;">\$--</div>	
CLAIMS	NUMBER FILED	NUMBER EXTRA	RATE		
Total claims	30 -20 =	10	X \$18.00	\$180	
Independent claims	3 -3 =	0	X \$80.00	\$0	
MULTIPLE DEPENDENT CLAIM(S) (if applicable)			+ 270.00	\$--	
TOTAL OF ABOVE CALCULATIONS =				\$1040	
<input checked="" type="checkbox"/> Applicant claims small entity status. See 37 CFR 1.27. The fees indicated above are reduced by 1/2.				\$520	
SUBTOTAL =				\$520	
Processing fee of \$130.00 for furnishing the English translation later than <input type="checkbox"/> 20 <input type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492(f)).				<div style="border: 1px solid black; padding: 5px; display: inline-block;">\$--</div>	
TOTAL NATIONAL FEE =				\$520	
Fee for recording the enclosed assignment (37 CFR 1.21(h)). The assignment must be accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31). \$40.00 per property +				<div style="border: 1px solid black; padding: 5px; display: inline-block;">\$--</div>	
TOTAL FEES ENCLOSED =				\$520	
				Amount to be:	\$
				refunded	\$
				charged	\$
a. <input checked="" type="checkbox"/> A check in the amount of \$520.00 to cover the above fees is enclosed.					
b. <input type="checkbox"/> Please charge my Deposit Account No. _____ in the amount of \$ _____ to cover the above fees. A duplicate copy of this sheet is enclosed.					
c. <input checked="" type="checkbox"/> The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any overpayment to Deposit Account No. 12-0080 . A duplicate copy of this sheet is enclosed.					
NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137(a) or (b)) must be filed and granted to restore the application to pending status.					
SEND ALL CORRESPONDENCE TO: Giulio A. DeConti, Esq. LAHIVE & COCKFIELD, LLP 28 State Street Boston, Massachusetts 02109 United States of America Date: 08 December 2000			<div style="font-size: 18pt; font-family: cursive; margin-bottom: 5px;">Megan E. Willams</div> <div style="border-bottom: 1px solid black; display: inline-block; width: 150px;"></div> SIGNATURE Megan E. Willams <hr style="border: 0; border-top: 1px solid black; margin: 5px 0;"/> NAME <div style="border-bottom: 1px solid black; display: inline-block; width: 100px; text-align: center;">43,270</div> REGISTRATION NUMBER		

09/719277

JC01 Rec'd PCT/PTO 08 DEC 2000

IN THE UNITED STATES PATENT DESIGNATED OFFICE (DO/US)
(National Phase of International App.: PCT/US99/12929)

In re the
application of: **Andrea D. BRANCH et al.**

International Application No.: **PCT/US99/12929**

International Filing Date: **09 JUNE 1999**

U.S. Serial No.: **Not yet assigned**

Filed: **Herewith**

For: **NOVEL HEPATITIS C VIRUS PEPTIDES
AND USES THEREOF**

Attorney Docket No.: **RII-003CPUS**

Assistant Commissioner for Patents
BOX PCT
Washington, D.C. 20231

Certification Under 37 CFR 1.10

I hereby certify that the attached:

Transmittal Letter (2 sheets in duplicate); Declaration, Petition and Power of Attorney (unexecuted) (4 sheets); PCT International Published Application (WO 99/63941) (without International Search Report) (51 sheets); Cover sheet of PCT International Published Application (WO 99/63941) (with International Search Report attached) (3 sheets); Sequence Listing (3 sheets); Diskette Containing Sequence Listing; Check (\$520) based on small entity; Certificate of Express Mailing (1 sheet); and Return Postcard are being deposited by me with the United States Postal Service "Express Mail Post Office to Addressee" service, Mailing Label No. **EL 589 754 944 US**, under 37 CFR 1.10 on the date indicated below and is addressed to the Assistant Commissioner for Patents, Box PCT, Washington, D.C. 20231.

Date: **08 December 2000**

Name: **Nelson F. Barros**

Signature: *Nelson Barros*

09/719277-0000

09/719277
JC01 Rec'd PCT/PTO 08 DEC 2000

SEQUENCE LISTING

<110> Andrea Branch, Jose Walewski, and Dechard Stump

5 <120> NOVEL HEPATITIS C VIRUS PEPTIDES AND USES THEREOF

<130> RII-003PC

10 <140>

<141>

<160> 8

15 <170> PatentIn Ver. 2.0

<210> 1

<211> 595

<212> DNA

20 <213> Hepatitis C virus

<400> 1

gcacgaatcc taaacctcaa agaaaaacca aacgtaacac caaccgtcgc ccacaggacg 60
tcaagttccc ggggtggcggc cagatcggtg gtggagttta cttgttgccg cgcagggggc 120
ctagattggg tgtgcgcgcg acgaggaaga cttccgagcg gtcgcaacct cgaggtagac 180
gtcagcctat ccccaaggca cgtcggcccg agggcaggac ctgggctcag cccgggtacc 240
cttggccctt ctatggcaat gaggggttgcg ggtgggcggg atggctcctg tctccccgtg 300
gctctcgccc tagctggggc cccacagacc cccggcgtag gtcgcgcaat ttgggtaagg 360
tcacgatac ccttacgtgc ggcttcgcgc acctcatggg gtacataccg ctcgtcggcg 420
ccctctttgg aggcgctgcc agggccctgg cgcattggcg ccgggttctg gaagacggcg 480
tgaactatgc aacagggaac ctctctgggt gctctttctc tatcttctt ctggccctgc 540
tctcttgctt gactgtgccg gcttcagcct accaagtgcg caattcctcg gggct 595

<210> 2

<211> 196

<212> PRT

<213> Hepatitis C virus

<400> 2

40 Ala Arg Ile Leu Asn Leu Lys Glu Lys Pro Asn Val Thr Pro Thr Val
1 5 10 15

Ala His Arg Thr Ser Ser Ser Arg Val Ala Val Arg Ser Leu Val Glu
20 25 30

45 Phe Thr Cys Cys Arg Ala Gly Ala Leu Asp Trp Val Cys Ala Arg Arg
35 40 45

50 Gly Arg Leu Pro Ser Gly Arg Asn Leu Glu Val Asp Val Ser Leu Ser
50 55 60

Pro Arg His Val Gly Pro Arg Ala Gly Pro Gly Leu Ser Pro Gly Thr
65 70 75 80

55 Leu Gly Pro Ser Met Ala Met Arg Val Ala Gly Gly Arg Asp Gly Ser
85 90 95

09/719277-0435
JC01 Rec'd PCT/PTO 08 DEC 2000

- 2 -

Cys Leu Pro Val Ala Leu Gly Leu Ala Gly Ala Pro Gln Thr Pro Gly
 100 105 110

5 Val Gly Arg Ala Ile Trp Val Arg Ser Ser Ile Pro Leu Arg Ala Ala
 115 120 125

Ser Pro Thr Ser Trp Gly Thr Tyr Arg Ser Ser Ala Pro Leu Leu Glu
 130 135 140

10 Ala Ala Pro Gly Pro Trp Arg Met Ala Ser Gly Phe Trp Lys Thr Ala
 145 150 155 160

Thr Met Gln Gln Gly Thr Phe Leu Val Ala Leu Ser Leu Ser Ser Phe
 165 170 175

15 Trp Pro Cys Ser Leu Ala Leu Cys Pro Leu Gln Pro Thr Lys Cys Ala
 180 185 190

20 Ile Pro Arg Gly
 195

<210> 3
 <211> 13
 25 <212> PRT
 <213> Hepatitis C virus

<220>
 <223> at location 8, X is N or K

<220>
 <223> at location 9, X is V or E

35 <220>
 <223> at location 13, X is A or V

<400> 3
 Leu Asn Leu Lys Glu Lys Pro Xaa Xaa Thr Pro Thr Xaa
 1 5 10

40 <210> 4
 <211> 13
 <212> PRT

45 <213> Hepatitis C virus

<220>
 <223> at location 6, X is L or S

50 <220>
 <223> at location 10, X is A or V

<220>
 <223> at location 11, X is A or V

55 <400> 4
 Ala Ala His Arg Thr Xaa Ser Ser Arg Xaa Xaa Val Arg
 1 5 10

- 3 -

<210> 5
<211> 14
5 <212> PRT
<213> Hepatitis C virus

<400> 5
10 Leu Asn Leu Lys Glu Lys Pro Asn Val Thr Pro Thr Ala Cys
1 5 10

<210> 6
<211> 14
15 <212> PRT
<213> Hepatitis C virus

<400> 6
20 Ala Ala His Arg Thr Ser Ser Ser Arg Ala Val Val Arg Cys
1 5 10

<210> 7
<211> 18
25 <212> PRT
<213> Hepatitis C virus

<400> 7
30 Pro Thr Asp Pro Arg Arg Arg Ser Arg Asn Leu Gly Lys Val Ile Asp
1 5 10 15
Thr Cys

<210> 8
<211> 20
35 <212> PRT
<213> Hepatitis C virus

<400> 8
40 Gly Cys Ala Thr Arg Lys Thr Ser Glu Arg Ser Gln Pro Arg Gly Arg
1 5 10 15

45 Arg Ala Pro Ile
20

CRF Errors Corrected by the STIC Systems Branch

Serial Number: 09/719,277

CRF Processing Date: 1/30/2001
 Edited by: A
 Verified by: A (STIC staff)

ENTERED

- ☐ Changed a file from non-ASCII to ASCII
- ☐ Changed the margins in cases where the sequence text was "wrapped" down to the next line.
- ☐ Edited a format error in the Current Application Data section, specifically: _____
- ☐ Edited the Current Application Data section with the actual current number. The number inputted by the applicant was ☐ the prior application data; or ☐ other _____
- ☐ Added the mandatory heading and subheadings for "Current Application Data".
- ☐ Edited the "Number of Sequences" field. The applicant spelled out a number instead of using an integer.
- ☐ Changed the spelling of a mandatory field (the headings or subheadings), specifically: _____
- ☐ Corrected the SEQ ID NO when obviously incorrect. The sequence numbers that were edited were: _____
- ☐ Inserted or corrected a nucleic number at the end of a nucleic line. SEQ ID NO's edited: _____
- ☐ Corrected subheading placement. All responses must be on the same line as each subheading. If the applicant placed a response below the subheading, this was moved to its appropriate place.
- ☐ Inserted colons after headings/subheadings. Headings edited included _____
- ☐ Deleted extra, invalid, headings used by an applicant, specifically: _____
- ☒ Deleted ☒ non-ASCII "garbage" at the beginning/end of files; ☐ secretary initials/filename at end of file;
☐ page numbers throughout text; ☐ other invalid text, such as _____
- ☐ Inserted mandatory headings, specifically: _____
- ☐ Corrected an obvious error in the response, specifically: _____
- ☐ Edited identifiers where upper case is used but lower case is required, or vice versa.
- ☐ Corrected an error in the Number of Sequences field, specifically: _____
- ☐ A "Hard Page Break" code was inserted by the applicant. All occurrences had to be deleted
- ☐ Deleted *ending* stop codon in amino acid sequences and adjusted the "(A)Length:" field accordingly (error due to a PatentIn bug). Sequences corrected: _____
- ☐ Other: _____

Examiner: The above corrections must be communicated to the applicant in the first Office Action. DO NOT send a copy of this form. 3/1/95

RAW SEQUENCE LISTING

DATE: 02/06/2001

PATENT APPLICATION: US/09/719,277

TIME: 10:28:41

Input Set : A:\Pto.amc

Output Set: N:\CRF3\02062001\I719277.raw

3 <110> APPLICANT: Andrea Branch, Jose Walewski, and Dechard Stump
 5 <120> TITLE OF INVENTION: NOVEL HEPATITIS C VIRUS PEPTIDES AND USES THEREOF
 8 <130> FILE REFERENCE: RII-003PC

C--> 10 <140> CURRENT APPLICATION NUMBER: US/09/719,277

C--> 11 <141> CURRENT FILING DATE: 2000-12-08

13 <160> NUMBER OF SEQ ID NOS: 8
 15 <170> SOFTWARE: PatentIn Ver. 2.0
 17 <210> SEQ ID NO: 1

18 <211> LENGTH: 595

19 <212> TYPE: DNA

20 <213> ORGANISM: Hepatitis C virus

22 <400> SEQUENCE: 1

23 gcacgaatcc taaacctcaa agaaaaacca aacgtaacac caaccgtcgc ccacaggagc 60
 24 tcaagttccc ggggtggcgt cagatcgttg gtggagttaa cttgttgccg cgcaggggccc 120
 25 ctgattggg tgtgcgcgcg acgaggaaga cttccgagcg gtgcgaacct cgaggtagac 180
 26 gtcagcctat ccccaaggca cgtcggcccg agggcaggac ctgggctcag cccgggtacc 240
 27 cttggcccct ctatggcaat gaggggtgcg ggtgggcggg atggctcctg tctcccctg 300
 28 gctctcggcc tagctggggc cccacagacc cccggcgtag gtgcgcgaat ttgggtaagg 360
 29 tcatcgatac ccttacgtgc ggtctgcgcg acctcatggg gtacataacc ctcgtcgcgc 420
 30 cccctcttgg aggcgctgcc agggccctgg cgcctggcgt ccgggttctg gaagacggcg 480
 31 tgaactatgc aacaggggaa cttcctgggt gctctttctc tatcttctt ctggccctgc 540
 32 tctcttgccct gactgtgccc gcttcagcct accaagtgcg caattcctcg gggct 595

34 <210> SEQ ID NO: 2

35 <211> LENGTH: 196

36 <212> TYPE: PRT

37 <213> ORGANISM: Hepatitis C virus

39 <400> SEQUENCE: 2

40 Ala Arg Ile Leu Asn Leu Lys Glu Lys Pro Asn Val Thr Pro Thr Val
 41 1 5 10 15
 43 Ala His Arg Thr Ser Ser Ser Arg Val Ala Val Arg Ser Leu Val Glu
 44 20 25 30
 46 Phe Thr Cys Cys Arg Ala Gly Ala Leu Asp Trp Val Cys Ala Arg Arg
 47 35 40 45
 49 Gly Arg Leu Pro Ser Gly Arg Asn Leu Glu Val Asp Val Ser Leu Ser
 50 50 55 60
 52 Pro Arg His Val Gly Pro Arg Ala Gly Pro Gly Leu Ser Pro Gly Thr
 53 65 70 75 80
 55 Leu Gly Pro Ser Met Ala Met Arg Val Ala Gly Gly Arg Asp Gly Ser
 56 85 90 95
 59 Cys Leu Pro Val Ala Leu Gly Leu Ala Gly Ala Pro Gln Thr Pro Gly
 60 100 105 110
 62 Val Gly Arg Ala Ile Trp Val Arg Ser Ser Ile Pro Leu Arg Ala Ala
 63 115 120 125
 65 Ser Pro Thr Ser Trp Gly Thr Tyr Arg Ser Ser Ala Pro Leu Leu Glu
 66 130 135 140
 68 Ala Ala Pro Gly Pro Trp Arg Met Ala Ser Gly Phe Trp Lys Thr Ala
 69 145 150 155 160

RAW SEQUENCE LISTING

DATE: 02/06/2001

PATENT APPLICATION: US/09/719,277

TIME: 10:28.41

Input Set : A:\Pto.amc

Output Set: N:\CRF3\02062001\I719277.raw

```

71 Thr Met Gln Gln Gly Thr Phe Leu Val Ala Leu Ser Leu Ser Ser Phe
72                               165                               170                               175
74 Trp Pro Cys Ser Leu Ala Leu Cys Pro Leu Gln Pro Thr Lys Cys Ala
75                               180                               185                               190
77 Ile Pro Arg Gly
78                               195

```

81 <210> SEQ ID NO: 3

82 <211> LENGTH: 13

83 <212> TYPE: PRT

84 <213> ORGANISM: Hepatitis C virus

86 <220> FEATURE:

87 <223> OTHER INFORMATION: at location 8, X is N or K

89 <220> FEATURE:

90 <223> OTHER INFORMATION: at location 9, X is V or E

92 <220> FEATURE:

93 <223> OTHER INFORMATION: at location 13, X is A or V

95 <400> SEQUENCE: 3

w06 96 Leu Asn Leu Lys Glu Lys Pro Xaa Xaa Thr Pro Thr Xaa
97 1 5 10

100 <210> SEQ ID NO. 4

101 <211> LENGTH: 13

102 <212> TYPE: PRT

103 <213> ORGANISM: Hepatitis C virus

105 <220> FEATURE:

106 <223> OTHER INFORMATION: at location 6, X is L or S

108 <220> FEATURE:

109 <223> OTHER INFORMATION: at location 10, X is A or V

111 <220> FEATURE:

112 <223> OTHER INFORMATION: at location 11, X is A or V

114 <400> SEQUENCE: 4

w06 115 Ala Ala His Arg Thr Xaa Ser Ser Arg Xaa Xaa Val Arg
116 1 5 10

119 <210> SEQ ID NO. 5

120 <211> LENGTH: 14

121 <212> TYPE: PRT

122 <213> ORGANISM: Hepatitis C virus

124 <400> SEQUENCE: 5

125 Leu Asn Leu Lys Glu Lys Pro Asn Val Thr Pro Thr Ala Cys

126 1 5 10

129 <210> SEQ ID NO: 6

130 <211> LENGTH: 14

131 <212> TYPE: PRT

132 <213> ORGANISM: Hepatitis C virus

134 <400> SEQUENCE: 6

135 Ala Ala His Arg Thr Ser Ser Ser Arg Ala Val Val Arg Cys

136 1 5 10

139 <210> SEQ ID NO: 7

140 <211> LENGTH: 18

141 <212> TYPE: PRT

Input Set : A:\Pto.amc
Output Set: N:\CRF3\02062001\I719277.raw

```

142 <213> ORGANISM: Hepatitis C virus
144 <400> SEQUENCE: 7
145 Pro Thr Asp Pro Arg Arg Arg Ser Arg Asn Leu Gly Lys Val Ile Asp
146   1                               5               10               15
148 Thr Cys
152 <210> SEQ ID NO: 8
153 <211> LENGTH: 20
154 <212> TYPE: PRT
155 <213> ORGANISM: Hepatitis C virus
157 <400> SEQUENCE: 8
158 Gly Cys Ala Thr Arg Lys Thr Ser Glu Arg Ser Gln Pro Arg Gly Arg
159   1                               5               10               15
161 Arg Ala Pro Ile
162           20

```

VERIFICATION SUMMARY

DATE: 02/06/2001

PATENT APPLICATION: US/09/719,277

TIME: 10 28:42

Input Set : A:\Pto.amc

Output Set: N:\CRF3\02062001\I719277.raw

L:10 M:270 C: Current Application Number differs, Replaced Application Number
L:11 M:271 C: Current Filing Date differs, Replaced Current Filing Date
L:96 M:258 W: Mandatory Feature missing, <221> not found for SEQ ID#:3
L:96 M:258 W: Mandatory Feature missing, <222> not found for SEQ ID#:3
L:96 M:340 W: (46) "n" or "Xaa" used: Feature required, for SEQ ID#:3
L:115 M:258 W: Mandatory Feature missing, <221> not found for SEQ ID#:4
L:115 M:258 W: Mandatory Feature missing, <222> not found for SEQ ID#:4
L:115 M:340 W: (46) "n" or "Xaa" used: Feature required, for SEQ ID#:4

F02F40/2267260

Does Not Comply
Corrected Diskette Needed

ERRORED SEQUENCES

Figure 1 consists of 12 histograms arranged in a 6x2 grid. The left column contains histograms for $n = 10, 20, 30, 40, 50, 60$. The right column contains histograms for $n = 70, 80, 90, 100, 110, 120$. Each histogram plots the 'Number of non-zero elements' on the x-axis against the 'Frequency' on the y-axis. The distributions are approximately bell-shaped and centered around $n/2$. The x-axis scale varies by histogram, and the y-axis scale also varies. The histograms show that as n increases, the distribution of non-zero elements becomes more concentrated around $n/2$.

VERIFICATION SUMMARY DATE 01/30/2001
 PATENT APPLICATION US/09/719,277 TIME 09:22 54

Input Set : A:\seqlist.txt
 Output Set: N:\CRF3\01302001\I719277.raw

L:10 M:270 C: Current Application Number differs, Replaced Application Number
 L:11 M:271 C: Current Filing Date differs, Replaced Current Filing Date
 L:96 M:258 W: Mandatory Feature missing, <221> not found for SEQ ID# 3
 L:96 M:258 W: Mandatory Feature missing, <222> not found for SEQ ID# 3
 L:96 M:340 W: (46) "n" or "Xaa" used. Feature required, for SEQ ID# 3
 L:115 M:258 W: Mandatory Feature missing, <221> not found for SEQ ID# 4
 L:115 M:258 W: Mandatory Feature missing, <222> not found for SEQ ID# 4
 L:115 M:340 W: (46) "n" or "Xaa" used. Feature required, for SEQ ID# 4
 L:164 M:333 E: Wrong sequence grouping, Amino acids not in groups'
 L:164 M:320 E: (1) Wrong Nucleic Acid Designator, NUMBER OF INVALID KEYS=1
 L:167 M:332 E: (32) Invalid/Missing Amino Acid Numbering, SEQ ID 8
 L:167 M:252 E: No. of Seq. differs. <211>LENGTH:Input:20 Found:21 SEQ 8

09494460

NOVEL HEPATITIS C VIRUS PEPTIDES AND USES THEREOFGovernment Funding

This work was funded, in part, by NIH grants DK 50795 and DK 52071. The
5 government may, therefore, have certain rights to this invention.

Background of the Invention

Hepatitis C virus (HCV) is closely related to both the pestivirus and flavivirus
genera in the *Flaviviridae* family. HCV is a single stranded RNA virus; the viral
10 genome is approximately 9.5 kb. HCV RNA is positive sense and has a unique open
reading frame which encodes a single polyprotein (Clarke. 1997. *J. Gen. Virol.*
78:2397). The polyprotein is proteolytically processed to yield the mature viral proteins
which include: nucleocapsid, envelope 1, envelope 2, metalloprotease, serine protease,
RNA helicase, cofactor, and RNA polymerase.

15 HCV is a major human pathogen. The virus was found to be the cause of most
cases of hepatitis which could not be ascribed to hepatitis A, hepatitis B, or hepatitis
delta virus (Clarke, *supra*). Over fifty percent of patients with hepatitis C virus (HCV)
become chronic carriers of the virus; there may be as many as 500 million chronic
carriers worldwide (Dhillon and Ducheiko. 1995. *Histopathology* 26: 297). Persistent
20 infection with the virus causes chronic hepatitis and may ultimately lead to cirrhosis
and/or cancer (Kuo et al. 1989. *Science* 244:362). Current therapies for HCV are
ineffective, consequently there is a need for new approaches to treat HCV infection.

Summary

25 The present invention is an important advance in the battle against hepatitis C.
The novel peptides of the invention, which are not encoded by the standard, polyprotein
HCV reading frame, have been shown to elicit an immune response in patients infected
with HCV and, thus, are produced during HCV infection. Accordingly, the invention
provides novel HCV polypeptides which are not derived from the HCV polyprotein and
30 methods of their use.

- 2 -

In one aspect, the invention pertains to an isolated or recombinant polypeptide or fragment thereof encoded by a nucleic acid molecule derived from a hepatitis C virus, which polypeptide has at least one of the following characteristics:

- 1) at least a portion of the polypeptide is encoded by a reading frame +1 or +2
5 relative to the standard hepatitis C virus open reading frame;
- 2) at least a portion of the polypeptide is encoded by a reading frame corresponding to the reading frame of SEQ ID NO:1 in which the first nucleotide of SEQ ID NO:1 is the first nucleotide of a codon;
- 3) at least a portion of the polypeptide comprises an amino acid sequence at least
10 60% identical to the amino acid sequence shown in SEQ ID NO:2; and
- 4) at least a portion of the polypeptide comprises an amino acid sequence encoded by a nucleic acid molecule which hybridizes under high stringency to the nucleotide sequence shown in SEQ ID NO:1.

In certain embodiments of the invention the novel HCV polypeptides or portion
15 thereof of claim 1 are at least about 8 amino acids to at least about 100 amino acids in length. In other embodiments, the polypeptides or portions thereof are at least about 14 amino acids to at least about 30 amino acids in length.

In one embodiment, the novel HCV polypeptides or portions thereof are encoded by a reading frame +1 or +2 to the standard hepatitis C reading frame. In preferred
20 embodiments, the polypeptides are encoded by a reading frame corresponding to the reading frame of SEQ ID NO:1 in which the first nucleotide of SEQ ID NO:1 is the first nucleotide of a codon. In other preferred embodiments, the polypeptides are encoded, in part by the nucleic acid molecule of SEQ ID NO:1 and cause an immune response in a subject.

In other embodiments, the novel HCV polypeptides comprise an amino acid
25 sequence at least 60% identical to the amino acid sequence shown in SEQ ID NO:2 and cause an immune response in a subject. In preferred embodiments, the novel HCV polypeptides comprise an amino acid sequence at least 90% identical to the amino acid sequence shown in SEQ ID NO:2 and causes an immune response in a subject. In other
30 preferred embodiments, the novel HCV polypeptides comprise an amino acid sequence shown in SEQ ID NO: 2 and cause an immune response in a subject.

In other embodiments, the novel HCV polypeptides comprise an amino acid sequence encoded by a nucleic acid molecule which hybridizes under high stringency to the nucleotide sequence shown in SEQ ID NO:1.

In other embodiments, the novel HCV polypeptides comprise at least a portion of
 5 an amino acid sequence selected from the group consisting of SEQ ID NO: 3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, and SEQ ID NO:8 and cause an immune response in a subject.

In still other embodiments, the invention pertains to isolated or recombinant polypeptides comprising an amino acid sequence selected from the group consisting of:
 10 LNLKEKP(X1)(X2)TPT(X3) and AAHRT(X4)SSR(X5)(X6)VR, wherein X1 is N or K, X2 is V or E, X3 is A or V, X4 is L or S, X5 is A or V, and X6 is A or V. In yet other embodiments, the novel HCV polypeptides consist of an amino acid sequence selected from the group consisting of LNLKEKPNVTPTAC and AAHRTSSRAVVRC.

In another aspect, the invention pertains to a vaccine composition for preventing
 15 hepatitis C infection in a subject. In one embodiment such a vaccine comprises a novel HCV polypeptide. In another embodiment, such a vaccine comprises a nucleic acid encoding a novel HCV polypeptide.

In another aspect, the invention pertains to an antibody which binds to a novel HCV polypeptide.
 20 In yet another aspect the invention pertains to a kit for detecting a hepatitis C infection. In one embodiment, such a kit comprises a novel HCV polypeptide. In another aspect, the kit comprises an antibody which binds to a novel HCV polypeptide.

In yet another aspect, the invention pertains to a method of preventing HCV infection by administering a novel HCV polypeptide to a subject or by causing said
 25 polypeptide to be synthesized in a subject prior to HCV infection such that HCV infection is prevented.

The invention also pertains to methods of diagnosing HCV infection. In one embodiment, the method comprises detecting the presence or absence of antibodies which react with a novel HCV polypeptide in the body fluid of a subject, wherein the
 30 presence of antibodies which bind the polypeptide is indicative of an infection with HCV. In another embodiment, the method comprises detecting the presence or absence

- 4 -

of a novel HCV polypeptide in the body fluid or tissue of a subject, wherein the presence of an HCV polypeptide is indicative of an infection with HCV.

In still another aspect, the invention pertains to a method for identifying a compound which interacts with a novel HCV polypeptide by contacting the polypeptide with a compound in a cell-free system under conditions which allow interaction of the compound with the polypeptide such that a complex is formed; separating the compounds which do not form complexes with an HCV polypeptide from those which do form complexes with an HCV polypeptide; and isolating and identifying the compounds which form complexes with an HCV polypeptide to identify a compound which interacts with a novel HCV polypeptide.

Detailed Description

The present invention is an important step forward in preventing Hepatitis C (HCV) infection, in treating ongoing infection, and in improving existing diagnostic techniques. The invention is based, in part, on the identification of novel polypeptides encoded by the Hepatitis C viral genome. These novel polypeptides are not encoded by the standard HCV polyprotein reading frame.

Before further description of the invention, certain terms employed in the specification, examples and appended claims are, for convenience, collected here.

I. Definitions

As used herein, the language "isolated or recombinant polypeptide" includes a polypeptide which is substantially free of cellular material or culture medium when produced by recombinant DNA techniques, or chemical precursors or other chemicals when chemically synthesized.

As used herein, the term "polypeptide or fragment thereof" includes full-length polypeptide molecules (from the first amino acid of translation initiation to the last amino acid prior to translation termination) and peptide portions of such molecules. Preferably the novel HCV polypeptides or fragments thereof are at least about 8 amino acids to at least about 100 amino acids in length. More preferably the polypeptides or fragments thereof are at least about 14 amino acids to at least about 30 amino acids in

- 5 -

length. In preferred embodiments, the novel HCV polypeptides of the invention comprise an amino acid sequence which is conserved among different HCV isolates. Such a conserved sequence can readily be determined using an alignment such as that provided in Table 1. In other embodiments, the novel HCV polypeptides comprise a

5 portion of a novel HCV amino acid sequence which is distal (carboxy terminal) to a stop codon in the +1 reading frame (relative to the main ORF) that includes the "UG" of the "AUG" that is the initiator codon of the main ORF. In a more preferred embodiment, the polypeptides or fragments thereof cause an immune response in a subject.

As used herein, the language "nucleic acid molecule" is intended to include DNA

10 molecules (e.g., cDNA or genomic DNA) and RNA molecules (e.g., genomic viral RNA or mRNA). The nucleic acid molecule may be single-stranded or double-stranded.

As used herein, the language "the standard hepatitis C virus open reading frame" is the open reading frame (ORF) of the viral RNA which encodes the well-known HCV polyprotein. The standard ORF represents the largest ORF in the viral genome. In the

15 infectious clone (GenBank accession number AF011751) the standard ORF uses nucleotide 342 as the first nucleotide of a codon and continues until nucleotide 9377. In different HCV isolates, the nucleotide which is a first nucleotide of a codon of the standard ORF may be at a slightly different position. The nucleotide which is a first nucleotide of a codon for any isolate can be easily be obtained to yield the standard HCV

20 ORF. For example, in the case of known isolates GenBank (or another database containing the nucleotide sequence information for the isolate) can be accessed and the coding sequence (CDS) information can be obtained. Alternatively, to determine the standard ORF of a known or a new isolate, the nucleic acid sequence of the known or new isolate can be aligned with a known sequence to give the highest homology (e.g.,

25 using a program such as BLAST). An exemplary BLAST search can be done, e.g., using the sequence found in GenBank accession number AF011751, as the query sequence. In this search, nucleotides 342-940 of AF011751 were used to search the non-redundant sequence database. The ORF of other HCV isolates which corresponds to the standard HCV ORF of AF011751 (in which the initiation codon is at position 342,

30 which is read as position 1 of the query sequence) can be read from the BLAST alignment. For example, the corresponding first nucleotide of a codon for GenBank

accession no. HPCCGAA is 342. Another way to find the standard ORF would be to use a program, such as Edit Seq. (DNASTAR) which is designed to identify ORFS using the AUG aligned with position 342 of AF011751 as the start codon.

The term "percent (%) identity" as used in the context of nucleotide and amino acid sequences (e.g., when one amino acid sequence is said to be X% identical to another amino acid sequence) refers to the percentage of identical residues shared between the two sequences, when optimally aligned. To determine the percent identity of two nucleotide or amino acid sequences, the sequences are aligned for optimal comparison purposes (e.g., gaps may be introduced in one sequence for optimal alignment with the other sequence). The residues at corresponding positions are then compared and when a position in one sequence is occupied by the same residue as the corresponding position in the other sequence, then the molecules are identical at that position. The percent identity between two sequences, therefore, is a function of the number of identical positions shared by two sequences (*i.e.*, % identity = # of identical positions/total # of positions x 100).

Computer algorithms known in the art can be used to optimally align and compare two nucleotide or amino acid sequences to define the percent identity between the two sequences. A preferred, non-limiting example of a mathematical algorithm utilized for the comparison of two sequences is the algorithm of Karlin and Altschul (1990) Proc. Natl. Acad. Sci. USA 87:2264-68, modified as in Karlin and Altschul (1993) Proc. Natl. Acad. Sci. USA 90:5873-77. Such an algorithm is incorporated into the NBLAST and XBLAST programs of Altschul, et al. (1990) J. Mol. Biol. 215:403-10. To obtain gapped alignments for comparison purposes, Gapped BLAST can be utilized as described in Altschul et al., (1997) Nucleic Acids Research 25(17):3389-3402. When utilizing BLAST and Gapped BLAST programs, the default parameters of the respective programs (e.g., XBLAST and NBLAST) can be used. See <http://www.ncbi.nlm.nih.gov>.

Another preferred, non-limiting example of a mathematical algorithm utilized for the comparison of sequences is the algorithm of Myers and Miller, CABIOS (1989). Such an algorithm is incorporated into the ALIGN program (version 2.0) which is part of the GCG sequence alignment software package. When utilizing the ALIGN program for

- 7 -

comparing amino acid sequences, a PAM120 weight residue table, a gap length penalty of 12, and a gap penalty of 4 can be used. If multiple programs are used to compare sequences, the program that provides optimal alignment (i.e., the highest percent identity between the two sequences) is used for comparison purposes.

5 As used herein, the language "+1 or +2 relative to the standard hepatitis C virus open reading frame" includes reading frames in which a first nucleotide of a codon is shifted +1 nucleotide relative to the standard ORF or +2 nucleotide relative to the standard ORF. The reading frames encoding the novel polypeptides do not necessarily contain an in-frame start codon.

10 As used herein, the language "the reading frame of SEQ ID NO:1" means that the first three nucleotides of the sequence shown in SEQ ID NO:1 are the first second and third nucleotides of a codon for translation into an amino acid of a polypeptide. The reading frame of SEQ ID NO:1 is +1 relative to the standard HCV ORF. The language "a reading frame corresponding to the reading frame of SEQ ID NO:1" means that when
15 a sequence from an HCV isolate other than the AF011751 isolate shown in SEQ ID NO:1 is aligned with the sequence of SEQ ID NO:1 to give the highest homology, e.g., using the BLAST program, it is then read in the same reading frame as SEQ ID NO:1 to give the reading frame corresponding the reading frame of SEQ ID NO:1. The nucleotide position of a first nucleotide of a codon of an HCV isolate which corresponds
20 to that of SEQ ID NO:1 may vary from isolate to isolate. An exemplary BLAST search illustrating this principle is provided as Appendix B. For example, for GenBank accession number AF009606 a first nucleotide of a codon in a reading frame which corresponds to the reading frame of SEQ ID NO:1 is nucleotide 346.

As used herein, the term "hybridizes under high stringency" is intended to
25 describe conditions for hybridization and washing under which nucleotide sequences at least 70 % homologous to each other typically remain hybridized to each other. Preferably, the conditions are such that sequences at least 75 %, 85%, or 95% identical to each other typically remain hybridized to each other. Such stringent conditions are known to those skilled in the art and can be found in *Current Protocols in Molecular*
30 *Biology*, John Wiley & Sons, N.Y. (1989), 6.3.1-6.3.6. A preferred, non-limiting example of stringent hybridization conditions are hybridization in 6X sodium

- 8 -

chloride/sodium citrate (SSC) at about 45°C, followed by one or more washes in 0.2 X SSC, 0.1% SDS at 50-65°C.

As used herein, the term "antibody" is intended to include immunoglobulin molecules and immunologically active portions of immunoglobulin molecules, *i.e.*,
5 molecules that contain an antigen binding site which specifically binds (immunoreacts with) an antigen, such as Fab and F(ab')₂ fragments. The terms "monoclonal antibodies" and "monoclonal antibody composition", as used herein, refer to a population of antibody molecules that contain only one species of an antigen binding site capable of immunoreacting with a particular epitope of an antigen, whereas the term "polyclonal
10 antibodies" and "polyclonal antibody composition" refer to a population of antibody molecules that contain multiple species of antigen binding sites capable of interacting with a particular antigen. A monoclonal antibody compositions thus typically display a single binding affinity for a particular antigen with which it immunoreacts.

As used herein, the term "adjuvant" includes agents which potentiate the immune
15 response to an antigen. Adjuvants can be administered in conjunction with the subject polypeptides to additionally augment the immune response.

As used herein, the term "enhancing an immune response" includes increasing T and/or B cell responses, *i.e.*, cellular and/or humoral immune responses, by treatment of a subject using the claimed methods. In one embodiment, the claimed methods can be
20 used to enhance T helper cell responses. In another embodiment, the claimed methods can be used to enhance cytotoxic T cell responses. The claimed methods can be used to enhance both primary and secondary immune responses. Preferably, the immune response is increased as compared to the response of immune cells to the antigen in the absence of treatment with the claimed methods. The immune response of a subject can
25 be determined by, for example, assaying antibody production, immune cell proliferation, the release of cytokines, the expression of cell surface markers, cytotoxicity, enhanced ability to clear infection with HCV, etc.

II. Novel HCV Polypeptides

30 The novel HCV polypeptides of the invention are not derived from an HCV polypeptide, *i.e.*, the polypeptides of the present invention are not encoded by the

09719277.041301

- 9 -

standard HCV ORF. These alternate reading frame polypeptides are translated from (or synthesized based on) a reading frame which is +1 or +2 to the standard HCV ORF. The position of the first nucleotide of an ORF in which these polypeptides are translated will vary slightly depending upon the isolate studied. For example, for the infectious clone

5 (GenBank accession number AF011751) the first nucleotide of the ORF in which the novel HCV polypeptides are translated is nucleotide 346, which is +5 relative to the standard HCV ORF. The first nucleotide of a codon of other, known or new isolates which results in a reading frame which corresponds to the reading frame of SEQ ID NO:1 can be determined, e.g., by performing a BLAST search using the nucleic acid

10 sequence of SEQ ID NO:1 as the query sequence as described above.

Translation of the novel HCV polypeptides of the invention does not necessarily have to begin at a start AUG codon. For example, previous work has shown that the start AUG of HCV could be mutated to AUU or CUG with little effect on translation efficiency (Clarke, supra). Alternatively, RNA editing may be involved in generating an

15 initiator codon. Translation of the novel HCV polypeptides may also begin at the initiation site of the standard HCV ORF with a frame shift into a different reading frame. Finally, translation may be initiated 5' of the AUG start codon of the standard ORF in any of the three reading frames, but shifted into the +1 reading frame (relative to the standard ORF) so as to yield production of peptides which are, in part, at least 60%

20 identical to a portion of SEQ ID NO. 2. The internal ribosome entry site (IRES) is a complex RNA structural element that includes part of the 5' untranslated region of HCV RNA and part of the adjacent coding region. It may induce frame shifting or translational by passing.

In preferred embodiments, the polypeptides are encoded by a reading frame

25 corresponding to the reading frame of SEQ ID NO:1 in which the first nucleotide of SEQ ID NO:1 is the first nucleotide of a codon. This reading frame can encode a polypeptide of at least about 126 amino acids in length before a termination codon is reached. Table 1 presents an alignment of novel HCV polypeptides which are encoded in this reading frame from various HCV isolates, along with a majority sequence derived

30 using the Clustal method of sequence alignment. Stop codons appear in certain of the isolates after amino acid 126. However, translation may proceed beyond these stop

FOIEB0112267260

- 10 -

codons. For example, in certain cases, these stop codons may be sequencing errors. Alternatively, readthrough can occur by mutation, altered transcription, RNA editing, frame shifting or ribosome slippage. Therefore, even in the polypeptides in which a stop codon appears, in certain embodiments of the invention, the novel HCV polypeptide
5 may be longer, i.e., translation may proceed past a termination codon. Therefore, in the case of, e.g., the infectious clone AF011751, translation of the polypeptide could terminate, for example, at position 163 or 186 of SEQ ID NO:2. When the novel HCV polypeptides of the invention are synthesized, these stop codons may be ignored.

In other embodiments, the polypeptides of the invention have some percentage
10 identity to the sequence shown in SEQ ID NO:2. The percent identity between two nucleic acid or amino acid sequences can easily be calculated by dividing the number of identical bases or amino acids by the total number of bases or amino acids. Sequences are aligned to give the highest percent identity and yet provide an alignment which is biologically meaningful. Sequences can be aligned manually or, preferably, using an
15 algorithm. For example, in the case of amino acid sequences, a FASTA search can be performed of the Swiss Protein database using the Biosum50.Cmp (scoring matrix). The gap creation penalty can be set, e.g., at 12 and the extension penalty can be set e.g., at 2. The joining threshold can be set, e.g., at 36; the optimization threshold can be set, e.g., at 24; and the optimization width can be set, e.g., at 16.

20 In certain embodiments, the novel HCV polypeptides comprise an amino acid sequence at least about 40-50% identical to the amino acid sequence shown in SEQ ID NO:2. In preferred embodiments, the novel HCV polypeptides comprise an amino acid sequence at least about 50-60% identical to the amino acid sequence shown in SEQ ID NO:2. In other preferred embodiments, the novel HCV polypeptides comprise an amino
25 acid sequence at least about 60-70% identical to the amino acid sequence shown in SEQ ID NO:2. In more preferred embodiments, the novel HCV polypeptides comprise an amino acid sequence at least about 70-80% identical to the amino acid sequence shown in SEQ ID NO:2. In still more preferred embodiments, the novel HCV polypeptides comprise an amino acid sequence at least about 80-90% identical to the amino acid
30 sequence shown in SEQ ID NO:2. In another preferred embodiment, the novel HCV polypeptides comprise an amino acid sequence shown in SEQ ID NO:2.

09719277 041301
PCT/US99/12929

In preferred embodiments, the polypeptides of the invention have the described percent identity over a length of at least about 10 amino acids. In more preferred embodiments, the percent identity of the polypeptides extends over a length of at least about 20-30 amino acids. In more preferred embodiment, the percent identity of the polypeptides extends over a length of at least about 30-40 amino acids. In a more preferred embodiment, the percent identity of the polypeptides extends over a length of at least about 40-50 amino acids. In another more preferred embodiment, the percent identity of the polypeptides extends over a length of more than 50 amino acids. In other preferred embodiments, the percent identity of the polypeptides extends over a length of more than 100 amino acids.

In other embodiments, the novel HCV polypeptides comprise an amino acid sequence encoded by a nucleic acid molecule having some percentage identity to the nucleic acid molecule shown in SEQ ID NO:1. In certain embodiments, the novel HCV polypeptides comprise an amino acid sequence encoded by a nucleic acid molecule at least 70% identical shown in SEQ ID NO:1 in which the polypeptide is encoded by the reading frame shown in SEQ ID NO:1. In preferred embodiments, the novel HCV polypeptides comprises an amino acid sequence encoded by a nucleic acid molecule at least 80% identical shown in SEQ ID NO:1 in which the polypeptide is encoded by the reading frame shown in SEQ ID NO:1. In more preferred embodiments, the novel HCV polypeptides comprise an amino acid sequence encoded by a nucleic acid molecule at least 90% identical shown in SEQ ID NO:1 in which the polypeptide is encoded by the reading frame shown in SEQ ID NO:1. In other embodiments, novel HCV polypeptides comprise an amino acid sequence encoded by a nucleic acid molecule shown in SEQ ID NO:1 in which the polypeptide is encoded by the reading frame shown in SEQ ID NO:1.

In certain embodiments, the novel HCV polypeptides of the invention are encoded by a nucleic acid molecule which hybridizes under stringent conditions to the nucleic acid sequence shown in SEQ ID NO:1. Stringent hybridization conditions are known in the art. In preferred embodiments, such polypeptides are encoded by a nucleic acid molecule which hybridizes under stringent conditions to a nucleic acid molecule from any HCV isolate, but which are read in or synthesized as if read in the reading frame of SEQ ID NO: 1.

In other embodiments, the novel HCV polypeptides comprise at least a portion of an amino acid sequence selected from the group consisting of SEQ ID NO: 3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, and SEQ ID NO:8 and cause an immune response in a subject. Other novel HCV polypeptides can be identified using an HCV nucleic acid sequence and determining the amino acids which are encoded in the +1 or +2 reading frame. Polypeptides of comprising these sequences can be made and assayed for reactivity with antibodies from infected subjects. Those polypeptides which bind to antibodies, i.e., have elicited an immune response in infected subjects are made by the virus during the course of infection and represent preferred novel HCV polypeptides.

In still other embodiments, the invention pertains to isolated or recombinant polypeptides comprising an amino acid sequence selected from the group consisting of: LNLKEKP(X1)(X2)TPT(X3) and AAHRT(X4)SSR(X5)(X6)VR, wherein X1 is N or K, X2 is V or E, X3 is A or V, X4 is L or S, X5 is A or V, and X6 is A or V. In yet other embodiments, the novel HCV polypeptides consist of an amino acid sequence selected from the group consisting of LNLKEKPNVTPTAC and AAHRTSSSRVAVRC.

In certain embodiments of the invention the novel HCV polypeptides of the invention are at least about 8 amino acids to at least about 100 amino acids in length. In other embodiments, the polypeptides of the invention are at least about 10 amino acids to at least about 50 amino acids in length. In other embodiments, the polypeptides of the invention are at least about 14 to at least about 25 amino acids in length.

In preferred embodiments, the novel HCV polypeptides are of a length sufficient to cause an immune response in a subject. Such an immune response can be measured using techniques which are known in the art. For example, the immune response elicited by the HCV polypeptides of the invention can be a T cell-mediated response which can be measured by, e.g., cytokine production and/or cellular proliferation or cellular cytotoxicity and/or a B cell mediated response which can be measured, e.g., by antibody production.

In certain embodiments the novel HCV polypeptides of the invention are made as fusion proteins. In addition to utilizing fusion proteins to enhance immunogenicity, fusion proteins can also facilitate the expression of proteins, including the novel HCV

polypeptides of the present invention. For example, a the novel HCV polypeptide can be generated as a glutathione-S-transferase (GST-fusion protein). Such GST-fusion proteins can enable easy purification of the novel HCV polypeptide, as for example by the use of glutathione-derivatized matrices (see, for example, *Current Protocols in Molecular Biology*, eds. Ausubel et al. (N.Y.: John Wiley & Sons, 1991)). In another embodiment, a fusion gene coding for a purification leader sequence, such as a poly-(His)/enterokinase cleavage site sequence, can be can be fused to a the novel HCV polypeptide, in order to permit purification of the poly(His)-the novel HCV polypeptide by affinity chromatography using a Ni²⁺ metal resin. The purification leader sequence can then be subsequently removed by treatment with enterokinase (e.g., see Hochuli et al. (1987) *J. Chromatography* 411:177; and Janknecht et al. *PNAS* 88:8972).

Techniques for making fusion genes are known to those skilled in the art. Essentially, the joining of various DNA fragments coding for different polypeptide sequences is performed in accordance with conventional techniques, employing blunt-ended or stagger-ended termini for ligation, restriction enzyme digestion to provide for appropriate termini, filling-in of cohesive ends as appropriate, alkaline phosphatase treatment to avoid undesirable joining, and enzymatic ligation. In another embodiment, the fusion gene can be synthesized by conventional techniques including automated DNA synthesizers. Alternatively, PCR amplification of gene fragments can be carried out using anchor primers which give rise to complementary overhangs between two consecutive gene fragments which can subsequently be annealed to generate a chimeric gene sequence (see, for example, *Current Protocols in Molecular Biology*, eds. Ausubel et al. John Wiley & Sons: 1992).

It will be understood that the preceding characteristics of HCV polypeptides are not mutually exclusive.

III. Production of Novel HCV Polypeptides

Novel HCV polypeptides can be produced by recombinant DNA techniques. For example, a nucleic acid molecule encoding such a polypeptide is cloned into an expression vector, the expression vector is introduced into a host cell and the novel HCV polypeptide is expressed in the host cell. The novel HCV polypeptide can then be

isolated from the cells by an appropriate purification scheme using standard protein purification techniques. As an alternative to recombinant expression, a novel HCV polypeptide can be synthesized chemically using standard peptide synthesis techniques or purchased commercially. Moreover, native novel HCV polypeptides can be isolated from cells (e.g., cultured human cells infected with HCV), for example using an antibody.

A. Recombinant Production of Novel HCV Polypeptides

In certain embodiments, the novel HCV polypeptides are encoded by a naturally-occurring HCV nucleic acid molecule. As used herein, a "naturally-occurring" nucleic acid molecule refers to an RNA molecule (or a DNA molecule derived therefrom) having a nucleotide sequence that occurs in nature (e.g., encodes a protein produced by a naturally occurring HCV isolate).

In addition to naturally-occurring isolates of the novel HCV polypeptides, the skilled artisan will further appreciate that changes may be introduced by mutation, e.g., into an HCV nucleotide sequence thereby leading to changes in the amino acid sequence of the encoded HCV polypeptides.

For example, an isolated nucleic acid molecule encoding a novel HCV polypeptide homologous to the polypeptide of SEQ ID NO: 2, i.e., having a certain percentage identity to the polypeptide of SEQ ID NO:2 can be created by introducing one or more nucleotide substitutions, additions or deletions into the nucleotide sequence of SEQ ID NO: 1 such that one or more amino acid substitutions, additions or deletions are introduced into the encoded polypeptide. Mutations can be introduced into SEQ ID NO: 1 by standard techniques, such as site-directed mutagenesis and PCR-mediated mutagenesis. Alternatively, such a polypeptide can be chemically synthesized to yield a polypeptide with a change in amino acid sequence from that in the naturally occurring polypeptide.

Preferably, no substitutions or conservative amino acid substitutions are made where there is high homology or identity in amino acid residues among the various isolates. A "conservative amino acid substitution" is one in which the amino acid residue is replaced with an amino acid residue having a similar side chain. Families of

- 15 -

amino acid residues having similar side chains have been defined in the art, including basic side chains (e.g., lysine, arginine, histidine), acidic side chains (e.g., aspartic acid, glutamic acid), uncharged polar side chains (e.g., glycine, asparagine, glutamine, serine, threonine, tyrosine, cysteine), nonpolar side chains (e.g., alanine, valine, leucine, isoleucine, proline, phenylalanine, methionine, tryptophan), beta-branched side chains (e.g., threonine, valine, isoleucine) and aromatic side chains (e.g., tyrosine, phenylalanine, tryptophan, histidine).

Alternatively, in another embodiment, mutations can be introduced randomly along all or part of a novel HCV polypeptide coding sequence, such as by saturation mutagenesis, and the resultant mutants can be screened, e.g., by testing for reactivity with antibodies from an individual with a past or present HCV infection.

B. Expression Vectors and Host Cells

The nucleic acid molecules described herein can be expressed in an expression vector to produce a novel HCV polypeptide. As used herein, the term "vector" refers to a nucleic acid molecule capable of transporting another nucleic acid to which it has been linked. One type of vector is a "plasmid", which refers to a circular double stranded DNA loop into which additional DNA segments may be ligated. Another type of vector is a viral vector, wherein additional DNA segments may be ligated into the viral genome. Certain vectors are capable of autonomous replication in a host cell into which they are introduced (e.g., bacterial vectors having a bacterial origin of replication and episomal mammalian vectors). Other vectors (e.g., non-episomal mammalian vectors) are integrated into the genome of a host cell upon introduction into the host cell, and thereby are replicated along with the host genome. Moreover, certain vectors are capable of directing the expression of genes to which they are operatively linked. Such vectors are referred to herein as "expression vectors". In general, expression vectors of utility in recombinant DNA techniques are often in the form of plasmids. In the present specification, "plasmid" and "vector" may be used interchangeably as the plasmid is the most commonly used form of vector. However, the invention is intended to include such other forms of expression vectors, such as viral vectors (e.g., replication defective

- 16 -

retroviruses, adenoviruses and adeno-associated viruses), which serve equivalent functions.

The recombinant expression vectors of the invention comprise a nucleic acid molecule as described herein in a form suitable for expression of the nucleic acid in a host cell, which means that the recombinant expression vectors include one or more regulatory sequences, selected on the basis of the host cells to be used for expression, which is operatively linked to the nucleic acid sequence to be expressed. Within a recombinant expression vector, "operably linked" is intended to mean that the nucleotide sequence of interest is linked to the regulatory sequence(s) in a manner which allows for expression of the nucleotide sequence (e.g., in an *in vitro* transcription/translation system or in a host cell when the vector is introduced into the host cell). The term "regulatory sequence" is intended to include promoters, enhancers and other expression control elements (e.g., polyadenylation signals). Such regulatory sequences are described, for example, in Goeddel; *Gene Expression Technology: Methods in Enzymology* 185, Academic Press, San Diego, CA (1990). Regulatory sequences include those which direct constitutive expression of a nucleotide sequence in many types of host cell and those which direct expression of the nucleotide sequence only in certain host cells (e.g., tissue-specific regulatory sequences). It will be appreciated by those skilled in the art that the design of the expression vector may depend on such factors as the choice of the host cell to be transformed, the level of expression of protein desired, etc. The expression vectors of the invention can be introduced into host cells to thereby produce novel HCV polypeptides, including fusion proteins comprising such polypeptides.

The recombinant expression vectors of the invention can be designed for expression of novel HCV polypeptides in prokaryotic or eukaryotic cells. For example, novel HCV polypeptides can be expressed in bacterial cells such as *E. coli*, insect cells (using baculovirus expression vectors) yeast cells or mammalian cells. Suitable host cells are discussed further in Goeddel, *Gene Expression Technology: Methods in Enzymology* 185, Academic Press, San Diego, CA (1990). Alternatively, the recombinant expression vector may be transcribed and translated *in vitro*, for example using T7 promoter regulatory sequences and T7 polymerase.

Expression of proteins in prokaryotes is most often carried out in *E. coli* with vectors containing constitutive or inducible promoters directing the expression of either fusion or non-fusion proteins. Fusion vectors add a number of amino acids to a protein encoded therein, usually to the amino terminus of the recombinant protein. Such fusion

5 vectors typically serve three purposes: 1) to increase expression of recombinant protein; 2) to increase the solubility of the recombinant protein; and 3) to aid in the purification of the recombinant protein by acting as a ligand in affinity purification. Often, in fusion expression vectors, a proteolytic cleavage site is introduced at the junction of the fusion moiety and the recombinant protein to enable separation of the recombinant protein from

10 the fusion moiety subsequent to purification of the fusion protein. Such enzymes, and their cognate recognition sequences, include Factor Xa, thrombin and enterokinase. Typical fusion expression vectors include pGEX (Pharmacia Biotech Inc; Smith, D.B. and Johnson, K.S. (1988) *Gene* 67:31-40), pMAL (New England Biolabs, Beverly, MA) and pRIT5 (Pharmacia, Piscataway, NJ) which fuse glutathione S-transferase (GST),

15 maltose E binding protein, or protein A, respectively, to the target recombinant protein.

Examples of suitable inducible non-fusion *E. coli* expression vectors include pTrc (Amann *et al.*, (1988) *Gene* 69:301-315) and pET 11d (Studier *et al.*, *Gene Expression Technology: Methods in Enzymology* 185, Academic Press, San Diego, California (1990) 60-89). Target gene expression from the pTrc vector relies on host

20 RNA polymerase transcription from a hybrid trp-lac fusion promoter. Target gene expression from the pET 11d vector relies on transcription from a T7 gn10-lac fusion promoter mediated by a coexpressed viral RNA polymerase (T7 gn1). This viral polymerase is supplied by host strains BL21(DE3) or HMS174(DE3) from a resident λ prophage harboring a T7 gn1 gene under the transcriptional control of the lacUV 5

25 promoter.

One strategy to maximize recombinant protein expression in *E. coli* is to express the protein in a host bacteria with an impaired capacity to proteolytically cleave the recombinant protein (Gottesman, S., *Gene Expression Technology: Methods in Enzymology* 185, Academic Press, San Diego, California (1990) 119-128). Another

30 strategy is to alter the nucleic acid sequence of the nucleic acid to be inserted into an expression vector so that the individual codons for each amino acid are those

preferentially utilized in *E. coli* (Wada et al., (1992) *Nuc. Acids Res.* 20:2111-2118).

Such alteration of nucleic acid sequences of the invention can be carried out by standard DNA synthesis techniques.

In another embodiment, the novel HCV polypeptides expression vector is a yeast
5 expression vector. Examples of vectors for expression in yeast *S. cerevisiae* include
pYepSec1 (Baldari. et al., (1987) *Embo J.* 6:229-234), pMFa (Kurjan and Herskowitz,
(1982) *Cell* 30:933-943), pJRY88 (Schultz et al., (1987) *Gene* 54:113-123), and pYES2
(Invitrogen Corporation, San Diego, CA).

Alternatively, novel HCV polypeptides can be expressed in insect cells using
10 baculovirus expression vectors. Baculovirus vectors available for expression of proteins
in cultured insect cells (e.g., Sf 9 cells) include the pAc series (Smith et al., (1983) *Mol.*
Cell Biol. 3:2156-2165) and the pVL series (Lucklow, V.A., and Summers, M.D., (1989)
Virology 170:31-39).

In yet another embodiment, a nucleic acid molecule encoding novel HCV
15 polypeptides of the invention is expressed in mammalian cells using a mammalian
expression vector. Examples of mammalian expression vectors include pCDM8 (Seed,
B., (1987) *Nature* 329:840) and pMT2PC (Kaufman et al. (1987), *EMBO J.* 6:187-195).
When used in mammalian cells, the expression vector's control functions are often
provided by viral regulatory elements. For example, commonly used promoters are
20 derived from polyoma, Adenovirus 2, cytomegalovirus and Simian Virus 40.

In another embodiment, the recombinant mammalian expression vector is
capable of directing expression of the nucleic acid preferentially in a particular cell type
(e.g., tissue-specific regulatory elements are used to express the nucleic acid). Tissue-
specific regulatory elements are known in the art. Non-limiting examples of suitable
25 tissue-specific promoters include the albumin promoter (liver-specific; Pinkert et al.
(1987) *Genes Dev.* 1:268-277), lymphoid-specific promoters (Calame and Eaton (1988)
Adv. Immunol. 43:235-275), in particular promoters of T cell receptors (Winoto and
Baltimore (1989) *EMBO J.* 8:729-733) and immunoglobulins (Banerji et al. (1983) *Cell*
33:729-740; Queen and Baltimore (1983) *Cell* 33:741-748), neuron-specific promoters
30 (e.g., the neurofilament promoter; Byrne and Ruddle (1989) *Proc. Natl. Acad. Sci. USA*
86:5473-5477), pancreas-specific promoters (Edlund et al. (1985) *Science* 230:912-916),

and mammary gland-specific promoters (e.g., milk whey promoter; U.S. Patent No. 4,873,316 and European Application Publication No. 264,166). Developmentally-regulated promoters are also encompassed, for example the murine hox promoters (Kessel and Gruss (1990) *Science* 249:374-379) and the α -fetoprotein promoter (Campes and Tilghman (1989) *Genes Dev.* 3:537-546).

A recombinant expression vector is introduced into a suitable host cell. The terms "host cell" and "recombinant host cell" are used interchangeably herein. It is understood that such terms refer not only to the particular subject cell but to the progeny or potential progeny of such a cell. Because certain modifications may occur in succeeding generations due to either mutation or environmental influences, such progeny may not, in fact, be identical to the parent cell, but are still included within the scope of the term as used herein.

A host cell may be any prokaryotic or eukaryotic cell. For example, novel HCV polypeptides may be expressed in bacterial cells such as *E. coli*, insect cells, yeast or mammalian cells (such as Chinese hamster ovary cells (CHO) or COS cells). Other suitable host cells are known to those skilled in the art.

Vector DNA can be introduced into prokaryotic or eukaryotic cells via conventional transformation or transfection techniques. As used herein, the terms "transformation" and "transfection" are intended to refer to a variety of art-recognized techniques for introducing foreign nucleic acid (e.g., DNA) into a host cell, including calcium phosphate or calcium chloride co-precipitation, DEAE-dextran-mediated transfection, lipofection, or electroporation. Suitable methods for transforming or transfecting host cells can be found in Sambrook et al. (*Molecular Cloning: A Laboratory Manual*, 2nd Edition, Cold Spring Harbor Laboratory press (1989)), and other laboratory manuals.

For stable transfection of mammalian cells, it is known that, depending upon the expression vector and transfection technique used, only a small fraction of cells may integrate the foreign DNA into their genome. In order to identify and select these integrants, a gene that encodes a selectable marker (e.g., resistance to antibiotics) is generally introduced into the host cells along with the gene of interest. Preferred selectable markers include those which confer resistance to drugs, such as G418,

- 20 -

hygromycin and methotrexate. Nucleic acid encoding a selectable marker may be introduced into a host cell on the same vector as that encoding the polypeptide or may be introduced on a separate vector. Cells stably transfected with the introduced nucleic acid can be identified by drug selection (e.g., cells that have incorporated the selectable
5 marker gene will survive, while the other cells die).

A host cell of the invention, such as a prokaryotic or eukaryotic host cell in culture, can be used to produce (i.e., express) novel HCV polypeptides. Accordingly, the invention further provides methods for producing novel HCV polypeptides using these host cells. In one embodiment, the method comprises culturing the host cell of
10 invention (into which a recombinant expression vector encoding novel HCV polypeptides has been introduced) in a suitable medium until a novel HCV polypeptide is produced. In another embodiment, the method further comprises isolating novel HCV polypeptides from the medium or the host cell.

15 ***C. Chemical Synthesis of Novel HCV Polypeptides***

The novel HCV polypeptides can be chemically synthesized as is well known in the art. Moreover, the peptide can be substituted and/or derivatized to optimize stability. The subject polypeptides can also be synthesized as branched polypeptides, particularly for vaccine applications as is known in the art (see, e.g., *Peptides*. Edited by Bernd Gutte
20 Academic Press 1995. pp. 456-493).

IV. Antibodies Which React With Novel HCV Polypeptides

In yet another aspect, the invention pertains to an antibody which binds to a novel HCV polypeptide. A novel HCV polypeptide, or fragment thereof, can be used as
25 an immunogen to generate antibodies that bind such a polypeptide using standard techniques for polyclonal and monoclonal antibody preparation. The invention provides numerous antigenic peptide fragments of novel HCV polypeptides for use as immunogens. Preferably, an antigenic peptide of such a polypeptide comprises at least 8 amino acid residues of the amino acid sequence shown in SEQ ID NO: 2 or the ARF #1
30 polypeptide or ARF #2 polypeptide consensus sequences. Preferably, the antigenic peptide comprises at least 10 amino acid residues, more preferably at least 14 amino acid

residues, even more preferably at least 18 amino acid residues. Preferred polypeptides comprise the ARF #1 consensus sequence :LNLKEKP(X1)(X2)TPT(X3) or the ARF#2 consensus sequence AAHRT(X4)SSR(X5)(X6)VR, wherein X1 is N or K, X2 is V or E, X3 is A or V, X4 is L or S, X5 is A or V, and X6 is A or V polypeptide sequences.

- 5 Other preferred HCV polypeptides comprise or consist of the sequence LNLKEKPNVTPTAC or AAHRTSSSRAVVR.

The subject HCV polypeptides are used to prepare antibodies by immunizing a suitable subject, (e.g., rabbit, goat, mouse or other mammal) with the immunogen. An appropriate immunogenic preparation can contain, for example, recombinantly expressed
10 novel HCV polypeptides or a chemically synthesized novel HCV polypeptide can be used. The preparation can further include an adjuvant, such as Freund's complete or incomplete adjuvant, or similar immunostimulatory agent. Immunization of a suitable subject with an immunogenic HCV polypeptides preparation induces a polyclonal HCV polypeptides antibody response.

- 15 Accordingly, another aspect of the invention pertains to antibodies which react with the novel HCV polypeptides. The term "antibody" as used herein refers to immunoglobulin molecules and immunologically active portions of immunoglobulin molecules, i.e., molecules that contain an antigen binding site which specifically binds (immunoreacts with) an HCV polypeptides. The invention provides polyclonal and
20 monoclonal antibodies that bind HCV polypeptides. The term "monoclonal antibody" or "monoclonal antibody composition", as used herein, refers to a population of antibody molecules that contain only one species of an antigen binding site capable of immunoreacting with a particular epitope of a novel HCV polypeptide. A monoclonal antibody composition thus typically displays a single binding affinity for a particular
25 HCV polypeptide with which it reacts.

Polyclonal anti-HCV polypeptide antibodies can be prepared as described above by immunizing a suitable subject with a HCV polypeptide immunogen or attenuated HCV virus, or can be obtained from an infected individual. The anti-HCV polypeptide antibody titer in the immunized subject can be monitored over time by standard
30 techniques, such as with an enzyme linked immunosorbent assay (ELISA) using immobilized HCV polypeptide. If desired, the antibody molecules directed against HCV

- polypeptide can be isolated from the animal (e.g., from the blood) and further purified by well known techniques, such as protein A chromatography to obtain the IgG fraction. At an appropriate time after immunization, e.g., when the antibody titers are highest, antibody-producing cells can be obtained from the subject and used to prepare
- 5 monoclonal antibodies by standard techniques, such as the hybridoma technique originally described by Kohler and Milstein (1975, *Nature* 256:495-497) (see also, Brown et al. (1981) *J. Immunol* 127:539-46; Brown et al. (1980) *J Biol Chem* 255:4980-83; Yeh et al. (1976) *PNAS* 76:2927-31; and Yeh et al. (1982) *Int. J. Cancer* 29:269-75), the more recent human B cell hybridoma technique (Kozbor et al. (1983) *Immunol*
- 10 *Today* 4:72), the EBV-hybridoma technique (Cole et al. (1985), *Monoclonal Antibodies and Cancer Therapy*, Alan R. Liss, Inc., pp. 77-96) or trioma techniques. The technology for producing monoclonal antibody hybridomas is well known (see generally R. H. Kenneth, in *Monoclonal Antibodies: A New Dimension In Biological Analyses*, Plenum Publishing Corp., New York, New York (1980); E. A. Lerner (1981) *Yale J.*
- 15 *Biol. Med.*, 54:387-402; M. L. Gefter et al. (1977) *Somatic Cell Genet.*, 3:231-36). Briefly, an immortal cell line (typically a myeloma) is fused to lymphocytes (typically splenocytes) from a mammal immunized with a HCV polypeptide immunogen as described above, and the culture supernatants of the resulting hybridoma cells are screened to identify a hybridoma producing a monoclonal antibody that binds the HCV
- 20 polypeptide.

- Any of the many well known protocols used for fusing lymphocytes and immortalized cell lines can be applied for the purpose of generating an anti-HCV polypeptide monoclonal antibody (see, e.g., G. Galfre et al. (1977) *Nature* 266:55052; Gefter et al. *Somatic Cell Genet.*, cited *supra*; Lerner, *Yale J. Biol. Med.*, cited *supra*;
- 25 Kenneth, *Monoclonal Antibodies*, cited *supra*). Moreover, the ordinary skilled worker will appreciate that there are many variations of such methods which also would be useful. Typically, the immortal cell line (e.g., a myeloma cell line) is derived from the same mammalian species as the lymphocytes. For example, murine hybridomas can be made by fusing lymphocytes from a mouse immunized with an immunogenic
- 30 preparation of the present invention with an immortalized mouse cell line. Preferred immortal cell lines are mouse myeloma cell lines that are sensitive to culture medium

containing hypoxanthine, aminopterin and thymidine ("HAT medium"). Any of a number of myeloma cell lines may be used as a fusion partner according to standard techniques, e.g., the P3-NS1/1-Ag4-1, P3-x63-Ag8.653 or Sp2/O-Ag14 myeloma lines. These myeloma lines are available from the American Type Culture Collection (ATCC),

- 5 Rockville, Md. Typically, HAT-sensitive mouse myeloma cells are fused to mouse splenocytes using polyethylene glycol ("PEG"). Hybridoma cells resulting from the fusion are then selected using HAT medium, which kills unfused and unproductively fused myeloma cells (unfused splenocytes die after several days because they are not transformed).

- 10 Hybridoma cells producing a monoclonal antibody of the invention are detected by screening the hybridoma culture supernatants for antibodies that bind HCV polypeptides, e.g., using a standard ELISA assay.

- Alternative to preparing monoclonal antibody-secreting hybridomas, a monoclonal anti-HCV polypeptide antibody can be identified and isolated by screening a
- 15 recombinant combinatorial immunoglobulin library (e.g., an antibody phage display library) with HCV polypeptides to thereby isolate immunoglobulin library members that bind HCV polypeptides. Kits for generating and screening phage display libraries are commercially available (e.g., the Pharmacia *Recombinant Phage Antibody System*, Catalog No. 27-9400-01; and the Stratagene *SurfZAPTM Phage Display Kit*, Catalog
- 20 No. 240612). Additionally, examples of methods and reagents particularly amenable for use in generating and screening antibody display library can be found in, for example, Ladner et al. U.S. Patent No. 5,223,409; Kang et al. International Publication No. WO 92/18619; Dower et al. International Publication No. WO 91/17271; Winter et al. International Publication WO 92/20791; Markland et al. International Publication No.
- 25 WO 92/15679; Breitling et al. International Publication WO 93/01288; McCafferty et al. International Publication No. WO 92/01047; Garrard et al. International Publication No. WO 92/09690; Ladner et al. International Publication No. WO 90/02809; Fuchs et al. (1991) *Bio/Technology* 9:1370-1372; Hay et al. (1992) *Hum Antibod Hybridomas* 3:81-85; Huse et al. (1989) *Science* 246:1275-1281; Griffiths et al. (1993) *EMBO J* 12:725-
- 30 734; Hawkins et al. (1992) *J Mol Biol* 226:889-896; Clarkson et al. (1991) *Nature* 352:624-628; Gram et al. (1992) *PNAS* 89:3576-3580; Garrad et al. (1991)

Bio/Technology 9:1373-1377; Hoogenboom et al. (1991) *Nuc Acid Res* 19:4133-4137; Barbas et al. (1991) *PNAS* 88:7978-7982; and McCafferty et al. *Nature* (1990) 348:552-554.

Additionally, recombinant anti-HCV polypeptide antibodies, such as chimeric
 5 and humanized monoclonal antibodies, comprising both human and non-human
 portions, which can be made using standard recombinant DNA techniques, are within
 the scope of the invention. Such chimeric and humanized monoclonal antibodies can be
 produced by recombinant DNA techniques known in the art, for example using methods
 described in Robinson et al. International Patent Publication PCT/US86/02269; Akira, et
 10 al. European Patent Application 184,187; Taniguchi, M., European Patent Application
 171,496; Morrison et al. European Patent Application 173,494; Neuberger et al. PCT
 Application WO 86/01533; Cabilly et al. U.S. Patent No. 4,816,567; Cabilly et al.
 European Patent Application 125,023; Better et al. (1988) *Science* 240:1041-1043; Liu
 et al. (1987) *PNAS* 84:3439-3443; Liu et al. (1987) *J. Immunol.* 139:3521-3526; Sun et
 15 al. (1987) *PNAS* 84:214-218; Nishimura et al. (1987) *Canc. Res.* 47:999-1005; Wood et
 al. (1985) *Nature* 314:446-449; and Shaw et al. (1988) *J. Natl Cancer Inst.* 80:1553-
 1559; Morrison, S. L. (1985) *Science* 229:1202-1207; Oi et al. (1986) *BioTechniques*
 4:214; Winter U.S. Patent 5,225,539; Jones et al. (1986) *Nature* 321:552-525;
 Verhoeyan et al. (1988) *Science* 239:1534; and Beidler et al. (1988) *J. Immunol.*
 20 141:4053-4060.

An anti-HCV polypeptide antibody (e.g., monoclonal antibody) can be used to
 isolate or detect HCV polypeptides by standard techniques, such as affinity
 chromatography, immunoprecipitation, ELISA, or RIA as is well known in the art. An
 anti-HCV polypeptide antibody can facilitate the purification of natural HCV
 25 polypeptide from cells and of recombinantly produced HCV polypeptides expressed in
 host cells. Moreover, an anti-HCV polypeptide antibody can be used to detect HCV
 polypeptides from a body fluid of a subject which is suspected to have an HCV
 infection. Detection may be facilitated by coupling (i.e., physically linking) the antibody
 to a detectable substance. Examples of detectable substances include various enzymes,
 30 prosthetic groups, fluorescent materials, luminescent materials and radioactive materials.
 Examples of suitable enzymes include horseradish peroxidase, alkaline phosphatase, β -

- 25 -

galactosidase, or acetylcholinesterase; examples of suitable prosthetic group complexes include streptavidin/biotin and avidin/biotin; examples of suitable fluorescent materials include umbelliferone, fluorescein, fluorescein isothiocyanate, rhodamine, dichlorotriazinylamine fluorescein, dansyl chloride or phycoerythrin; an example of a luminescent material includes luminol; and examples of suitable radioactive material include ^{125}I , ^{131}I , ^{35}S or ^3H .

Such antibodies can be incorporated in diagnostic kits and are also useful in passive immunization against HCV in patients which have an active HCV infection or are likely to be exposed to HCV.

10

IV. Uses of Novel HCV Polypeptides

In another aspect, the invention pertains to a vaccine composition which is administered to a subject prior to exposure to HCV to preventing hepatitis C infection in the subject. In one embodiment, the vaccine comprises a novel HCV polypeptide of the invention. In another embodiment, the vaccine causes a novel HCV polypeptide of the invention to be synthesized in a subject.

15

A. Vaccines

Novel HCV polypeptide sequences appropriate for use in vaccine compositions for the prevention of HCV in a subject can easily be determined. For example, epitopes which elicit an immune response can be identified by screening in an immunoassay against sera from patients with a past or ongoing HCV infection. Alternatively, immunogenic polypeptides can be identified by computer analysis to identify immunogenic epitopes. Finally, the full-length novel polypeptide could be used in a vaccine.

20

In another embodiment, agents which are known adjuvants can be administered with the subject polypeptides. At this time, the only adjuvant widely used in humans has been alum (aluminum phosphate or aluminum hydroxide). Saponin and its purified component Quil A, Freund's complete adjuvant and other adjuvants used in research and veterinary applications have potential use in human vaccines. However, new chemically defined preparations such as muramyl dipeptide, monophosphoryl lipid A,

30

FOI b7D " 4267260

phospholipid conjugates such as those described by Goodman-Snitkoff et al. *J. Immunol.* 147:410-415 (1991) resorcinols, non-ionic surfactants such as polyoxyethylene oleyl ether and n-hexadecyl polyethylene ether, enzyme inhibitors include pancreatic trypsin inhibitor, diisopropylfluorophosphate (DEP) and trasylol can also be used. In

5 embodiments in which antigen is administered, the antigen can e.g., be encapsulated within a proteoliposome as described by Miller et al., *J. Exp. Med.* 176:1739-1744 (1992) and incorporated by reference herein, or in lipid vesicles, such as Novasome™ lipid vesicles (Micro Vesicular Systems, Inc., Nashua, N. H.), to further enhance immune responses.

10 In yet other embodiments, as an alternative to administering the novel HCV polypeptide, the polypeptide can be synthesized by the subject. This can be done using a plasmid DNA construct which is similar to those used for delivery of reporter or therapeutic genes. Such a construct preferably comprises a bacterial origin of replication that allows amplification of large quantities of the plasmid DNA; a prokaryotic

15 selectable marker gene; a nucleic acid sequence encoding a novel HCV polypeptide or portion thereof; eukaryotic transcription regulatory elements to direct gene expression in the host cell; and a polyadenylation sequence to ensure appropriate termination of the expressed mRNA (Davis. 1997. *Curr. Opin. Biotechnol.* 8:635). Vectors used for DNA immunization may optionally comprise a signal sequence (Michel et al. 1995. *Proc.*

20 *Natl. Acad. Sci USA.* 92:5307; Donnelly et al. 1996. *J. Infect Dis.* 173:314). DNA vaccines can be administered by a variety of means, for example, by injection (e.g., intramuscular, intradermal, or the biolistic injection of DNA-coated gold particles into the epidermis with a gene gun that uses a particle accelerator or a compressed gas to inject the particles into the skin (Haynes et al. 1996. *J. Biotechnol.* 44:37)).

25 Alternatively, DNA vaccines can be administered by non-invasive means. For example, pure or lipid-formulated DNA can be delivered to the respiratory system or targeted elsewhere, e.g., Peyer's patches by oral delivery of DNA (Schubbert. 1997. *Proc. Natl. Acad. Sci. USA* 94:961). Attenuated microorganisms can be used for delivery to mucosal surfaces. (Sizemore et al. 1995. *Science.* 270:29)

Any of the instant vaccine compositions can comprise (or encode) one or more epitopes (either contiguous or non contiguous) of a novel HCV polypeptide. Such preparations can further comprise polypeptide sequences derived from an HCV polyprotein sequence. In other embodiments, such a vaccine composition can further

5 comprise a compound which will enhance the immunological reactivity of the novel HCV polypeptide epitope. For example, the immunogenicity of the novel HCV polypeptides may be enhanced by making a fusion proteins comprising a novel HCV polypeptide fused to a different polypeptide, i.e., not a novel HCV polypeptide. Techniques for making such fusion proteins are known in the art. Alternatively, a

10 vaccine can comprise an immunoregulatory molecule, such as a cytokine. For example, in one embodiment, plasmids for DNA vaccination can express a single immunogen, or two sequences can be coexpressed. In one embodiment, the additional sequences can be additional immunogens (novel HCV polypeptides or HCV polyprotein polypeptides or other polypeptides) or can encode modulators of immune responses such as lymphokine

15 genes or costimulatory molecules (Iwasaki et al. 1997. *J. Immunol.* 158:4591)

Typically, vaccine compositions of the present invention are prepared as injectables, either as liquid solutions or suspensions. Solid forms suitable for solution

in, or suspension in, liquid prior to injection may also be prepared. The composition may also be emulsified, or the polypeptide encapsulated into liposomes. The

20 polypeptide may be mixed with pharmaceutically acceptable excipients, for example, water, saline, dextrose, glycerol, ethanol, or the like. The composition may also comprise minor amounts of, for example, wetting agents, pH buffering agents and/or adjuvants, such as aluminum hydroxide, N-acetyl-muramyl-L-threonyl-D-isoglutamine (thr-MDP), N-acetyl-nor-muramyl-L-alanyl-D-isoglutamine (CGP 11637 or nor-MDP),

25 N-acetylmuramyl-L-alanyl-D-isoglutaminyl-L-alanine-2-(1'-2'-dipalmitoyl-sn-glycerol-3-hydroxyphosphoryloxy)ethylamine (CGP 19835A, or MTP-PE), or bacterial components.

Such vaccine compositions are generally administered parenterally, by injection, usually wither subcutaneously or intramuscularly. Other formulations may be

30 administered orally, by inhalation or as suppositories.

- 28 -

The polypeptides may be incorporated into the vaccine in a neutral or salt form. Pharmaceutically acceptable salts include the acid addition salts (formed with free amino groups of the polypeptide) and which are formed with inorganic acids, such as, for example, hydrochloric or phosphoric acids, or such organic acids such as acetic, oxalic, tartaric, maleic, or the like. Salts formed with the free carboxyl groups may also be derived from inorganic bases such as, for example, sodium, potassium, ammonium, calcium, or ferric hydroxides, and such organic bases as isopropylamine, trimethylamine, 2-ethylamino ethanol, histidine, procaine, or the like.

The vaccines are administered so as to be compatible with the dosage formulation, and in such an amount as will be prophylactically and/or therapeutically effective. The quantity to be administered depends on the subject to be treated, the capacity of the subject's immune system to mount an immune response to the vaccine, and the degree of protection desired. The range of 5 µg to 250 µg of antigen per dose, however, is often appropriate. The vaccine compositions may be given in a single dose or in multiple doses. Appropriate doses are well within the skill of the art to determine, and do not constitute undue experimentation.

In still another embodiment, the invention pertains to a method of preventing HCV in a subject by administering a novel HCV polypeptide to a subject or by causing a novel HCV polypeptide to be expressed in a subject.

B. Diagnostic Kits

In another aspect of the invention, methods for diagnosing HCV infection e.g., either a past or present infection, and diagnostic kits are provided for detecting an infection with HCV.

In one embodiment, the invention provides a method of diagnosing HCV infection by detecting the presence or absence of antibodies in the body fluid of a subject which bind to a novel HCV polypeptide. In one embodiment the method comprises incubating a test sample under conditions which allow the binding of a novel HCV polypeptide and an antibody in the test sample of body fluid and detecting the binding of polypeptide and antibody.

Test samples can be derived from any appropriate body fluid or tissue preparation, for example, whole blood, plasma, serum, spinal fluid, lymph fluid, tears, saliva, milk, or liver tissue preparations.

Detection of the binding between a novel HCV polypeptide of the invention and an antibody can be accomplished using any technique which is known in the art and can be facilitated using antibodies labelled as described above.

Antibodies which bind to novel HCV polypeptide can be detected using a number of different screening assays known in the art, such as an enzyme-linked immunosorbent assay (ELISA), a radioimmunoassay (RIA), or a Western Blot Assay. Each assay generally detects the presence of protein-antibody complexes of particular interest by employing a labeled reagent (e.g., an antibody) specific for the complex of interest. Accordingly, in the present invention, these assays are used to detect novel HCV polypeptide-antibody complexes formed between immunoglobulins (e.g., human IgG, IgM and IgA) contained in a biological sample and a novel HCV polypeptide. As will be described below, these protein-antibody complexes are preferably detected using an enzyme-linked antibody or antibody fragment (e.g., a monoclonal antibody or fragment thereof) which recognizes and specifically binds to the polypeptide-antibody complexes.

In one embodiment of the method, a sandwich ELISA assay is used. For example, a novel HCV polypeptide with or without conjugation to a carrier, such as activated BSA is immobilized on a plate. A body fluid sample from an individual is contacted with a novel HCV polypeptide under conditions which allow binding of the antibodies in the sample to the polypeptides. The sample is then removed, and any antibody which has bound to the HCV polypeptide is detected by contacting the sample with a labeled secondary antibody or antibody fragment which binds to an antibody which might be present in the subjects sample, e.g., an anti-human antibody. The unbound secondary antibody is removed and the presence of secondary antibody which remains bound is detected, e.g., using a label as described above. Possible controls for use in the method include body fluids from uninfected subjects and polypeptides which are not novel HCV polypeptides. In accordance with the present invention, the presence of such an antibody is indicative of an infection with HCV.

In another embodiment of the assay the test sample can be tested for the presence of novel HCV polypeptides using known antibodies. In these embodiments, antibodies that bind to novel HCV polypeptides are used to detect the presence of novel HCV polypeptides in the body fluid of a subject or in a cell of a subject. In performing
5 such an assay, the antibodies which bind to a novel HCV polypeptide are contacted with a cell or body fluid of a subject under conditions where a novel HCV polypeptide in the subject's sample can bind to the antibody. Unbound antibody is removed and bound antibody is detected. Any of the antibodies described above can be used in practicing this method. Preferred antibodies for use in the methods of this embodiment are highly
10 specific, including monospecific and, more preferably, monoclonal antibodies or fragments thereof. In preferred embodiments, such antibodies are labelled. In other embodiments, such antibodies are detected by employing a secondary antibody which binds to them and not to the test sample from a subject. The presence of a novel HCV polypeptide in the subject's sample is indicative of an infection with HCV.

15 In yet another aspect, the present invention provides an assay kit for diagnosing HCV infection in a subject. Preferably, the kit contains a solid support (e.g., an ELISA plate) capable of adsorbing immunoglobulin (e.g., IgG, IgM and IgA) from a subject's sample (preferably a human biological sample, such as a body fluid) and a monoclonal antibody or fragment thereof specific for a novel HCV polypeptide. In
20 another embodiment, the solid support can be omitted from the kit. In another embodiment, the kit contains a solid support (e.g., an ELISA plate or a slide) and a monoclonal antibody or fragment thereof specific for a novel HCV polypeptide. In other embodiments, the solid support can be omitted. The assay kit can optionally include instructions, or additional reagents such as a solution for washing unbound proteins from
25 the solid support, and materials needed for performing a detection assay.

C. Targets for Therapeutic Intervention

The novel HCV polypeptides of the invention are also targets for anti-HCV therapy. As such the invention provides methods for identifying compounds which
30 interact with a novel HCV polypeptide and, thus, are likely to interfere with infection. In one embodiment, the method involves contacting the polypeptide with a compound in

a cell-free system under conditions which allow interaction of the compound with the polypeptide such that a complex is formed. The complexes of polypeptide and compound can then be separated from the compounds which do not bind to the HCV polypeptide, the compounds which bind to HCV polypeptides can then be isolated and
5 identified.

Exemplary compounds which can be screened for activity in the subject assays include, but are not limited to, peptides, nucleic acids, carbohydrates, small organic molecules, and natural product extract libraries. The term "non-peptidic compound" is intended to encompass compounds that are comprised, at least in part, of molecular
10 structures different from naturally-occurring L-amino acid residues linked by natural peptide bonds. However, "non-peptidic compounds" are intended to include compounds composed, in whole or in part, of peptidomimetic structures, such as D-amino acids, non-naturally-occurring L-amino acids, modified peptide backbones and the like, as well
15 to naturally-occurring L-amino acid residues linked by natural peptide bonds. "Non-peptidic compounds" also are intended to include natural products.

A recent trend in medicinal chemistry includes the production of mixtures of compounds, referred to as libraries. While the use of libraries of peptides is well established in the art, new techniques have been developed which have allowed the
20 production of mixtures of other compounds, such as benzodiazepines (Bunin et al. 1992. J. Am. Chem. Soc. 114:10987; DeWitt et al. 1993. Proc. Natl. Acad. Sci. USA 90:6909) peptoids (Zuckermann. 1994. J. Med. Chem. 37:2678) oligocarbamates (Cho et al. 1993. Science. 261:1303), and hydantoins (DeWitt et al. supra). Rebek et al. have described an approach for the synthesis of molecular libraries of small organic molecules
25 with a diversity of 104-105 (Carell et al. 1994. Angew. Chem. Int. Ed. Engl. 33:2059; Carell et al. Angew. Chem. Int. Ed. Engl. 1994. 33:2061).

The compounds of the present invention can be obtained using any of the numerous approaches in combinatorial library methods known in the art, including: biological libraries; spatially addressable parallel solid phase or solution phase libraries,
30 synthetic library methods requiring deconvolution, the 'one-bead one-compound' library method, and synthetic library methods using affinity chromatography selection. The

biological library approach is limited to peptide libraries, while the other four approaches are applicable to peptide, non-peptide oligomer or small molecule libraries of compounds (Lam, K.S. *Anticancer Drug Des.* 1997. 12:145).

In one embodiment, the test compound is a peptide or peptidomimetic. In another, preferred embodiment, the compounds are small, organic non-peptidic compounds.

Other exemplary methods for the synthesis of molecular libraries can be found in the art, for example in: Erb et al. 1994. *Proc. Natl. Acad. Sci. USA* 91:11422; Horwell et al. 1996 *Immunopharmacology* 33:68; and in Gallop et al. 1994. *J. Med. Chem.* 37:1233.

Libraries of compounds may be presented in solution (e.g., Houghten (1992) *Biotechniques* 13:412-421), or on beads (Lam (1991) *Nature* 354:82-84), chips (Fodor (1993) *Nature* 364:555-556), bacteria (Ladner USP 5,223,409), spores (Ladner USP '409), plasmids (Cull et al. (1992) *Proc Natl Acad Sci USA* 89:1865-1869) or on phage (Scott and Smith (1990) *Science* 249:386-390); (Devlin (1990) *Science* 249:404-406); (Cwirla et al. (1990) *Proc. Natl. Acad. Sci.* 87:6378-6382); (Felici (1991) *J. Mol. Biol.* 222:301-310); (Ladner *supra.*).

In many drug screening programs which test libraries of compounds and natural extracts, high throughput assays are desirable in order to maximize the number of compounds surveyed in a given period of time. Assays which are performed in cell-free systems, such as may be derived with purified or semi-purified proteins, are often preferred as "primary" screens in that they can be generated to permit rapid development and relatively easy detection of an alteration in a molecular target which is mediated by a test compound. Accordingly, in an exemplary screening assay of the present invention, the compound of interest is contacted with a novel HCV polypeptide. Detection and quantification of novel HCV polypeptide/compound complexes identifies the compound as a potential modulator of a novel HCV polypeptide. The efficacy of the compound can be assessed by generating dose response curves from data obtained using various concentrations of the test compound. Moreover, a control, e.g., using a different polypeptide can also be performed to provide a baseline for comparison.

Complex formation between the novel HCV polypeptide and a compound may be detected by a variety of techniques. For instance, modulation of the formation of complexes can be quantitated using, for example, detectably labelled proteins such as radiolabelled, fluorescently labelled, or enzymatically labelled novel HCV polypeptides, by immunoassay, or by chromatographic detection.

Typically, it will be desirable to immobilize either the novel HCV polypeptide or the compound or both to facilitate separation of compound/novel HCV complexes from uncomplexed forms, as well as to accommodate automation of the assay. In one embodiment, a fusion protein can be provided which adds a domain that allows the polypeptide to be bound to a matrix. For example, glutathione-S-transferase/receptor (GST/receptor) fusion protein forms of the novel polypeptides can be adsorbed onto glutathione sepharose beads (Sigma Chemical, St. Louis, MO) or glutathione derivatized microtitre plates, which are then combined with the test compound which is bound to beads and incubated under conditions conducive to complex formation, e.g. at physiological conditions for salt and pH, though slightly more stringent conditions may be desired, e.g., at 4°C in a buffer containing 0.6M NaCl or a detergent such as 0.1% Triton X-100. Following incubation, the uncomplexed forms are removed by washing and compounds which bind to novel HCV polypeptides are identified.

Other techniques for immobilizing proteins on matrices are also available for use in the subject assay. For instance, the novel HCV polypeptides can be immobilized utilizing conjugation of biotin and streptavidin. For instance, biotinylated novel HCV polypeptides can be prepared from biotin-NHS (N-hydroxy-succinimide) using techniques well known in the art (e.g., biotinylation kit, Pierce Chemicals, Rockford, IL), and immobilized in the wells of streptavidin-coated 96 well plates (Pierce Chemical). Alternatively, antibodies reactive with the novel HCV polypeptides can be derivatized to the wells of the plate, and the novel HCV polypeptides trapped in the wells by antibody conjugation. As above, preparations of a novel HCV polypeptide and a test compound are incubated in the wells of the plate, and the amount of novel HCV polypeptides/ compound complex trapped in the well can be quantitated.

- Other exemplary methods for detecting such complexes, in addition to those described above for the GST-immobilized complexes, include immunodetection of complexes using antibodies reactive with the novel HCV polypeptide, or which are reactive with the receptor protein and compete for binding with the novel HCV
- 5 polypeptide; as well as enzyme-linked assays which rely on detecting an enzymatic activity associated with the novel HCV polypeptide. In the instance of the latter, the enzyme can be chemically conjugated or provided as a fusion protein with the novel HCV polypeptide. To illustrate, the novel HCV polypeptide can be chemically cross-linked or genetically fused with alkaline phosphatase, and the amount of novel HCV
- 10 polypeptide trapped in the complex can be assessed with a chromogenic substrate of the enzyme, e.g. paranitrophenylphosphate. Likewise, a fusion protein comprising the novel HCV polypeptide and glutathione-S-transferase can be provided, and complex formation quantitated by detecting the GST activity using 1-chloro-2,4-dinitrobenzene (Habig et al (1974) *J Biol Chem* 249:7130).
- 15 For processes which rely on immunodetection for quantitating one of the proteins trapped in the complex, antibodies against the protein, such as the anti-novel HCV antibodies described herein, can be used. Alternatively, the protein to be detected in the complex can be "epitope tagged" in the form of a fusion protein which includes, in addition to the novel HCV polypeptide, a second polypeptide for which antibodies are
- 20 readily available (e.g. from commercial sources). For instance, the GST fusion proteins described above can also be used for quantification of binding using antibodies against the GST moiety. Other useful epitope tags include myc-epitopes (e.g., see Ellison et al. (1991) *J Biol Chem* 266:21150-21157) which includes a 10-residue sequence from c-myc, as well as the pFLAG system (International Biotechnologies, Inc.) or the pEZZ-
- 25 protein A system (Pharmacia, NJ).
- The practice of the present invention will employ, unless otherwise indicated, conventional techniques of cell biology, cell culture, molecular biology, microbiology, recombinant DNA, and immunology, which are within the skill of the art. Such techniques are explained fully in the literature. See, for example, *Genetics; Molecular*
- 30 *Cloning A Laboratory Manual*, 2nd Ed., ed. by Sambrook, J. et al. (Cold Spring Harbor Laboratory Press (1989)); *Short Protocols in Molecular Biology*, 3rd Ed., ed. by

- Ausubel, F. *et al.* (Wiley, NY (1995)); *DNA Cloning*, Volumes I and II (D. N. Glover ed., 1985); *Oligonucleotide Synthesis* (M. J. Gait ed. (1984)); Mullis *et al.* U.S. Patent No: 4,683,195; *Nucleic Acid Hybridization* (B. D. Hames & S. J. Higgins eds. (1984)); the treatise, *Methods In Enzymology* (Academic Press, Inc., N.Y.); *Immunochemical*
- 5 *Methods In Cell And Molecular Biology* (Mayer and Walker, eds., Academic Press, London (1987)); *Handbook Of Experimental Immunology*, Volumes I-IV (D. M. Weir and C. C. Blackwell, eds. (1986)); and Miller, J. *Experiments in Molecular Genetics* (Cold Spring Harbor Press, Cold Spring Harbor, N.Y. (1972)).
- 10 The contents of all references, pending patent applications and published patents, cited throughout this application are hereby expressly incorporated by reference. Specifically, the contents of USSN 60/088,670, titled Novel Hepatitis C Virus Peptides And Uses Thereof, filed on June 9, 1998 and 60/089,138, titled Novel Hepatitis C Virus
- 15 Peptides And Uses Thereof, filed on June 11, 1998 are incorporated herein by this reference.

The invention is further illustrated by the following examples, which should not be construed as further limiting.

20 Examples

Example 1. The detection of antibodies against novel HCV polypeptides.

- Consensus polypeptides were synthesized based on the sequence homology
- 25 between novel HCV polypeptides shown in Table 1. The following peptides were made using conventional techniques by Biosynthesis Corp. (Lewisville, TX): LNLKEKPNVTPTAC and AAHRTSSSRVAVRC (alternate reading frame (ARF) polypeptides 1 and 2, respectively). The following polypeptides derived from HCV CORE protein were used as controls: PTDPRRRSRNLGKVIDTC and
- 30 GCATRKTSESRQPRGRRAPI. The peptides were conjugated to activated bovine serum albumin giving up to 4 peptide molecules for each BSA molecule. Peptide-BSA

- 36 -

conjugates at a concentration of approximately 0.5 mg/ml were shipped on ice in 3 mls of phosphate buffered saline, 0.1 M, pH 7.4. They were aliquoted and stored at -20°C until use.

ELISAs were performed as described by Kirkegaard and Perry Laboratories, Inc. (Gaithersburg, MD). For use in the ELISA, the polypeptides were dissolved in 1X "coating buffer" to give a peptide-BSA concentration of 1 µg/ml; 100 µl were added to 96 well microtiter plates (Falcon 3072, Becton Dickinson and Company, Franklin Lakes, NJ) and incubated at room temperature for 1 hour or overnight at 4°C. The coating solution was removed and plates were then blocked by adding 300 µl of 1% BSA in phosphate buffered saline (as prepared by Kirkegaard and Perry), and incubating for 30 min. at room temperature. The blocking solution was removed and 100 µl of 1% BSA (1X) was added.

Sera was obtained from patients which were known to have or to have previously had an HCV infection. Serum samples were diluted 1:100 in 1X BSA and 100 µl was added to the first well of each (yielding a total of 200 µl). Serum samples were then serially diluted (two-fold at each step). Control wells contained BSA only. Plates were incubated for 1 hour at room temperature with moderate agitation to allow binding. Plates were washed 5 times with 1X wash solution (PBS and 0.02% Tween). After washing, 100 µl of the secondary antibody was immediately added and allowed to react for 1 hour at room temperature. The secondary antibody was either the Fab fragment of anti-human IgG conjugated to horse radish peroxidase (HRP) at a dilution of 1:1000, or anti-human IgG (in PBS with 1% BSA). Plates were washed 5 times with 1X washing solution, and then 100 µl of hydrogen peroxide and TMB were added and allowed to react for 10 to 30 minutes at room temperature. To stop the reaction, 100 µl of 1M phosphoric acid were added. O.D. measurements were obtained by using a dual wavelength scanner: 450 nm values-650 nm values.

Results of ELISA Tests for Antibodies of HCV-Specific Polypeptides

30

Control Sera

HCV Patient Sera

- 37 -

	Mean OD (S.D.) (N=2, tested in duplicate)	Mean OD (S.D.) (N=6, tested in duplicate)
CORE #1	0.379 (0.063)	0.851 (0.231)
5 CORE #2	0.775 (0.053)	1.390 (0.400)
ARF #1	0.763 (0.133)	2.286 (0.215)
ARF #2	0.401 (0.039)	0.813 (0.813)

Example 2. Development of a western blot assay for detecting antibody production to
 10 alternate HCV reading frame proteins.

Five micrograms of BSA-alternate reading frame peptide conjugates and appropriate controls (quenched BSA, BSA-BSA peptide conjugates) were denatured by incubating in standard Lamelli loading buffer (with beta-mercaptoethanol
 15 and SDS), at 100°C for three minutes. Samples were then cooled and spun in a microcentrifuge prior to loading on a discontinuous 1.5 mm thick SDS-PAGE gel (4% stacking gel [pH 6.8] and 12.5% [pH 8.4] resolving gel). The running buffer was TRIS-glycine/SDS (0.1%), with an unadjusted pH of approximately 8.4.

Samples were run through the stacking gel (1 cm) for approximately 45 mins at
 20 90 volts. After the bromophenol blue (BB) entered the resolving gel, the voltage was increased to 160 V. The gels were run for approximately 1.5 hrs, or until the BB was 2/3 of the way through the gel. After stopping the run, the gels were either stained with coomassie blue, or equilibrated for 15 minutes with the transfer buffer (1X CAPS, pH 11.0, 10% MeOH. PVDF membranes (Immobilon-P
 25 [0.45 micron pore size], were wet in 100% MeOH for 1 min, then 50:50 MeOH:double distilled water for 5 mins, and finally equilibrated with the transfer buffer. The transfer was set up in transfer buffer (filter paper, gel, PVDF membrane, and filter paper), and slipped into the BIO-RAD transfer tank. The transfer was run at 20 volts, overnight (approximately 10 hrs), at 4°C, with stirring.
 30 After transfer, the tank was disassembled, the membrane was soaked in 100% MeOH for one minute, then allowed to air-dry. In order to visualize the

0971927, 044304

- 38 -

efficiency of the transfer, the membranes (after re-wetting in MeOH then water) were incubated with 1% Ponceau-S red stain. and rinsed in double distilled water. After scanning the image, the dye was removed in a dilute NaOH solution (1ml saturated NaOH in 100mls ddwater.

- 5 The membranes were then blocked in 3% NFDM (6 grams non-fat dried milk in 200mls of 1x TBS [TRIS-buffered saline, ph 7.4), for one hour. After blocking, the membranes were rinsed in two washes of 1xTBS, 200 mls apiece.

- The membranes were then incubated with the primary antibody solution 4 mls of a 1/200 dilution of patient sera in 1% NFDM in 1XTTBS (TBS with 0.025% Tween-20). This incubation lasted one hour, at 30oC, in glass with slow rotation of the tube or beaker. After this incubation, the membranes were washed three times in 200 mls of 1X TTBS.
- 10

- The secondary antibody solution was 200 mls of a 1/3000 dilution of the BIO-RAD goat anti-human alkaline phosphatase conjugate, in 1% NFDM in 1X TTBS. This incubation lasted 1 hr, at 30oC, with gentle shaking.
- 15

 After this incubation, the membranes were washed twice with 200 ml of 1X TTBS (5 mins apiece), and one with 200 mls of 1X TBS.

- The bands were visualized with the BIO-RAD AP substrate kit (200 mls total), with gentle, occasional shaking. After visualization, the membranes were washed several times with ddwater, followed by one wash with MeOH, then air-dried and photographed or scanned.
- 20

T06140 4267260

TABLE 1. Novel HCV Polypeptides

A Q I L N L K E K P N V T P T A A H R T L S S R V A V R S L A E F T C C R A G A P D W V C A R L	Majority
A R I L N L K E K P N V T P T V A H R T S S S R V A V R S L V E F T C C R A G A L D W V C A R R	AF011751
A H F L N L K E K P K E T P S V A H R T S S S R V A D R S L V E Y T C C R A G A H D W V C A R R	D17763
A Q I L N L K E K P K E T Q T A A H R T L S S R V A V R S L A E F T C C R A G A P G W V C A R Q	D10988
A R I L N L K E K P N V T P T A A H R T L S S R V A A R S L A E F T C C R A G A P E W V C A R R	D14853
A Q I L N L K E K P K E T P T V A H K T L S F R A A R S L A E Y T C C R A G A P G W V C A R Q	D00944
A Q I Q N P K D K P K E T P T V A H R T S S S R A V V R S W V E Y T C C R A G A L D W V C A R L	D63822
A R I L N L K E K P N V T P T A A Q W T L S S R V V A R S L A E F T C C R A G A P D W V C A R L	Y11604
A Q I L N L K E K P N V T P T A A H R T S S S R A V V R S L V E F T C C R A G A P G W V C A R L	D50482
G R L P S G R S L V E G A S L S P R I A G P R A G P G L S P G T L G P S M A M R V A G G Q D G S C P	Majority
G R L P S G R N L E V D V S L S P R H V C P R A G P G L S P G T L G P S M A M R V A G G R D G S C L	AF011751
V K L L N G H S L A D D D S L S P R R V G A K A G P G L S P G T L G P S M V T R A A G Q G G S C P	D17763
G R L L S D P S R V D D A S P S R K I G A P A S P G E S Q D I L G P C T E T R V A A G R V G S C P	D10988
G R L P S G R N L A G G V S L F P R P A D P R E G P G R S P G T L G P S M A T R A V G G R D P S C P	D14853
G R L R S G P S H V E G A S P S L R I G A P L A N P G E N Q D T P G P Y T G M R D S A G Q D R S C P	D00944
G R L P N G P S P E A G V S P F Q R L A A R R A V P G V S L G T H G P C M G M R A A G G Q G G S C P	D63822

1.036T.410" 226T.260

R11-003CPPC

G R L R S G R N L V E D A N L S P R R V D P R E G P G H N Q D I H G L F T V M R V V G G Q D G S C P
G R L P S G R N L V E G D N L S P R F A G P R A G P G L S P G T L G P S M A M R V W G G Q D G S C H

Y11604
D50482

WO 99/63941

4

P A A L G L L G A P M T P G G G P A I W V R S S I P L R A A S P T S W G T S R S S A P L - G A S P E
Majority
AF011751
D17763
D10988
D14853
D00944
D63822
Y11604
D50482

5

L W R M A 5 G S W R T G - I T Q Q G I F P V A L F L S S F W L F C P A - L S Q L R P S K C A T L V G
Majority
AF011751
D17763
D10988
D14853
D00944
D63822
Y11604
D50482

What is claimed is:

1. An isolated or recombinant polypeptide or fragment thereof encoded by a nucleic acid molecule derived from a hepatitis C virus, having at least one of the following
5 characteristics:
 - 1) at least a portion of the polypeptide is encoded by a reading frame +1 or +2 relative to the standard hepatitis C virus open reading frame;
 - 2) at least a portion of the polypeptide is encoded by a reading frame corresponding to the reading frame of SEQ ID NO:1 in which the first nucleotide of
10 SEQ ID NO:1 is the first nucleotide of a codon;
 - 3) at least a portion of the polypeptide comprises an amino acid sequence at least 60% identical to the amino acid sequence shown in SEQ ID NO:2; and
 - 4) at least a portion of the polypeptide comprises an amino acid sequence encoded by a nucleic acid molecule which hybridizes under high stringency to the
15 nucleotide sequence shown in SEQ ID NO:1.
2. The polypeptide or portion thereof of claim 1, wherein said polypeptide is at least about 8 amino acids to at least about 100 amino acids in length.
- 20 3. The polypeptide or portion thereof of claim 2, wherein said polypeptide is at least about 14 amino acids to at least about 30 amino acids in length.
4. The polypeptide or portion thereof of claim 1, wherein said polypeptide is encoded by a reading frame +1 or +2 to the standard hepatitis C reading frame.
- 25 5. The polypeptide or portion thereof of claim 1, wherein said polypeptide is encoded by a reading frame corresponding to the reading frame of SEQ ID NO:1 in which the first nucleotide of SEQ ID NO:1 is the first nucleotide of a codon.

0971927 041301
PCT/HO/226760

6. The polypeptide or portion thereof of claim 5, wherein said polypeptide or portion thereof is encoded by the nucleic acid molecule of SEQ ID NO:1 and causes an immune response in a subject.
- 5 7. The polypeptide or portion thereof of claim 1, wherein said polypeptide comprises an amino acid sequence at least 60% identical to the amino acid sequence shown in SEQ ID NO:2 and causes an immune response in a subject.
8. The polypeptide or portion thereof of claim 1, wherein said polypeptide
10 comprises an amino acid sequence at least 90% identical to the amino acid sequence shown in SEQ ID NO:2 and causes an immune response in a subject.
9. The polypeptide or portion thereof of claim 1, wherein said polypeptide
15 comprises an amino acid sequence shown in SEQ ID NO: 2 which polypeptide causes an immune response in a subject.
10. The polypeptide or portion thereof of claim 1, wherein said polypeptide
20 comprises an amino acid sequence encoded by a nucleic acid molecule which hybridizes under high stringency to the nucleotide sequence shown in SEQ ID NO:1.
11. The polypeptide or portion thereof of claim 1 which polypeptide comprises at least a portion of an amino acid sequence selected from the group consisting of SEQ ID NO: 3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, and SEQ ID NO:8 and causes an immune response in a subject.
- 25 12. An isolated or recombinant polypeptide comprising an amino acid sequence selected from the group consisting of: LNLKEKP(X1)(X2)TPT(X3) and AAHRT(X4)SSR(X5)(X6)VR, wherein X1 is N or K, X2 is V or E, X3 is A or V, X4 is L or S, X5 is A or V, and X6 is A or V.
- 30

09719277 041301

13. A polypeptide consisting of an amino acid sequence selected from the group consisting of LNLKEKPNVTPTAC and AAHRTSSSRVVRRC.
14. A vaccine composition for preventing hepatitis C infection in a subject comprising the polypeptide of claim 1.
15. A vaccine composition for preventing hepatitis C infection in a subject comprising the polypeptide of claim 2.
- 10 16. A vaccine composition for preventing hepatitis C infection in a subject comprising the polypeptide of claim 4.
17. A vaccine composition for preventing hepatitis C infection in a subject comprising the polypeptide of claim 7.
- 15 18. A vaccine composition for preventing hepatitis C infection in a subject comprising the polypeptide of claim 12.
19. A vaccine composition for preventing hepatitis C infection in a subject comprising a nucleic acid encoding polypeptide of claim 1.
- 20 20. A vaccine composition for preventing hepatitis C infection in a subject comprising a nucleic acid encoding polypeptide of claim 2.
- 25 21. A vaccine composition for preventing hepatitis C infection in a subject comprising a nucleic acid encoding polypeptide of claim 4.
22. A vaccine composition for preventing hepatitis C infection in a subject comprising a nucleic acid encoding polypeptide of claim 7.

23. A vaccine composition for preventing hepatitis C infection in a subject comprising a nucleic acid encoding polypeptide of claim 12.
24. An antibody which binds to a polypeptide of claim 1.
- 5 25. A kit for detecting a hepatitis C infection comprising the polypeptide of claim 1.
26. A kit for detecting a hepatitis C infection comprising an antibody to the polypeptide of claim 1.
- 10 27. A method of preventing HCV infection comprising administering the polypeptide of claim 1 to a subject or by causing said polypeptide to be synthesized in a subject prior to HCV infection such that HCV infection is prevented.
- 15 28. A method of diagnosing HCV infection comprising detecting the presence or absence of antibodies which react with the polypeptide of claim 1 in the body fluid of a subject, wherein the presence of antibodies which bind the polypeptide is indicative of an infection with HCV.
- 20 29. A method of diagnosing HCV infection comprising detecting the presence or absence of the polypeptide of claim 1 in the body fluid or tissue of a subject, wherein the presence of an HCV polypeptide is indicative of an infection with HCV.

30. A method for identifying a compound which interacts with the polypeptide of claim 1, comprising:

contacting said polypeptide with a compound in a cell-free system under conditions which allow interaction of the compound with the polypeptide such that a

5 complex is formed;

separating the compounds which do not form complexes with an HCV polypeptide from those which do form complexes with an HCV polypeptide; and

isolating and identifying the compounds which form complexes with an HCV polypeptide.

094927 044304
T06T40 225T460

- 1 -

SEQUENCE LISTING

<110> Andrea Branch, Jose Walewski, and Dechard Stump

5 <120> NOVEL HEPATITIS C VIRUS PEPTIDES AND USES THEREOF

<130> RII-003PC

10 <140>
<141>

<160> 8

15 <170> PatentIn Ver. 2.0

<210> 1
<211> 595
<212> DNA

20 <213> Hepatitis C virus

<400> 1
gcaacgaatcc taaacctcaa agaaaaacca aacgtaacac caaccgtcgc ccacaggacg 60
tcaagttccc ggggtggcggc cagatcggtg gtggagtta cttgttgccg cgcaggggcc 120
25 ctagattggg tgtgcgcgcg acgaggaaga cttccgagcg gtcgcaacct cgaggtagac 180
gtcagcctat ccccaaggca cgtcggcccg agggcaggac ctgggctcag cccgggtacc 240
cttggccoct ctatggcaat gagggttgcg ggtgggcggg atggctcctg tctccccgtg 300
gctctcggcc tagctggggc cccacagacc ccggcgtag gtcgcgcaat ttgggtaagg 360
tcatcgatac ccttacgtgc ggcttcgccg acctcatggg gtacataccg ctgcgcggcg 420
30 cccctcttgg aggcgctgcc agggccctgg cgcattggcg ccgggttctg gaagacggcg 480
tgaactatgc aacaggggac cttcctgggt gctctttctc tatcttcctt ctggccctgc 540
tctctgcct gactgtgccc gcttcagcct accaagtgcg caattcctcg gggct 595

<210> 2
35 <211> 196
<212> PRT
<213> Hepatitis C virus

<400> 2
40 Ala Arg Ile Leu Asn Leu Lys Glu Lys Pro Asn Val Thr Pro Thr Val
1 5 10 15

Ala His Arg Thr Ser Ser Ser Arg Val Ala Val Arg Ser Leu Val Glu
20 25 30

45 Phe Thr Cys Cys Arg Ala Gly Ala Leu Asp Trp Val Cys Ala Arg Arg
35 40 45

Gly Arg Leu Pro Ser Gly Arg Asn Leu Glu Val Asp Val Ser Leu Ser
50 55 60

Pro Arg His Val Gly Pro Arg Ala Gly Pro Gly Leu Ser Pro Gly Thr
65 70 75 80

55 Leu Gly Pro Ser Met Ala Met Arg Val Ala Gly Gly Arg Asp Gly Ser
85 90 95

PCT/US99/12929

- 2 -

Cys Leu Pro Val Ala Leu Gly Leu Ala Gly Ala Pro Gln Thr Pro Gly
 100 105 110

5 Val Gly Arg Ala Ile Trp Val Arg Ser Ser Ile Pro Leu Arg Ala Ala
 115 120 125

Ser Pro Thr Ser Trp Gly Thr Tyr Arg Ser Ser Ala Pro Leu Leu Glu
 130 135 140

10 Ala Ala Pro Gly Pro Trp Arg Met Ala Ser Gly Phe Trp Lys Thr Ala
 145 150 155 160

Thr Met Gln Gln Gly Thr Phe Leu Val Ala Leu Ser Leu Ser Ser Phe
 165 170 175

15 Trp Pro Cys Ser Leu Ala Leu Cys Pro Leu Gln Pro Thr Lys Cys Ala
 180 185 190

20 Ile Pro Arg Gly
 195

25 <210> 3
 <211> 13
 <212> PRT
 <213> Hepatitis C virus

30 <220>
 <223> at location 8, X is N or K

35 <220>
 <223> at location 9, X is V or E

35 <220>
 <223> at location 13, X is A or V

40 <400> 3
 Leu Asn Leu Lys Glu Lys Pro Xaa Xaa Thr Pro Thr Xaa
 1 5 10

45 <210> 4
 <211> 13
 <212> PRT
 <213> Hepatitis C virus

50 <220>
 <223> at location 6, X is L or S

50 <220>
 <223> at location 10, X is A or V

55 <220>
 <223> at location 11, X is A or V

55 <400> 4
 Ala Ala His Arg Thr Xaa Ser Ser Arg Xaa Xaa Val Arg
 1 5 10

00127.0460
 00127.0460
 00127.0460

- 3 -

<210> 5
 <211> 14
 5 <212> PRT
 <213> Hepatitis C virus

 <400> 5
 Leu Asn Leu Lys Glu Lys Pro Asn Val Thr Pro Thr Ala Cys
 10 1 5 10

 <210> 6
 <211> 14
 15 <212> PRT
 <213> Hepatitis C virus

 <400> 6
 Ala Ala His Arg Thr Ser Ser Ser Arg Ala Val Val Arg Cys
 20 1 5 10

 <210> 7
 <211> 18
 25 <212> PRT
 <213> Hepatitis C virus

 <400> 7
 Pro Thr Asp Pro Arg Arg Arg Ser Arg Asn Leu Gly Lys Val Ile Asp
 30 1 5 10 15
 Thr Cys

 35
 <210> 8
 <211> 20
 <212> PRT
 <213> Hepatitis C virus
 40
 <400> 8
 Gly Cys Ala Thr Arg Lys Thr Ser Glu Arg Ser Gln Pro Arg Gly Arg
 1 5 10 15
 45 Arg Ala Pro Ile
 20

0974997460
 04430

**DECLARATION, PETITION AND POWER OF ATTORNEY
FOR PATENT APPLICATION**

(Check one):

- ☐ Declaration Submitted with Initial Filing
☒ Declaration Submitted after Initial Filing

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name,

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:

NOVEL HEPATITIS C VIRUS PEPTIDES AND USES THEREOF

the specification of which (check one):

- ☐ is attached hereto.

OR

- ☒ was filed on 09 June 1999 as International Application Number
PCT/US99/12929

- ☐ and was amended by PCT Article 19 Amendment on _____
(if applicable),

- ☐ and was amended by PCT Article 34 Amendment on _____
(if applicable).

I acknowledge the duty to disclose to the Office all information known to me to be material to patentability as defined in Title 37, Code of Federal Regulations, §1.56.

I hereby state that I have reviewed and understood the contents of the above-identified specification, including the claims, as amended by any amendment referred to above.

091999-04301

PRIORITY CLAIM

(Check one):

- ☐ no such applications have been filed.
- ☒ such applications have been filed as follows

1) FOREIGN PRIORITY CLAIM: I hereby claim foreign priority benefits under Title 35, United States Code, §119(a)-(d) or §365(b) of any foreign application(s) for patent or inventor's certificate or §365(a) of any PCT international application which designated at least one country other than the United States of America, listed below and have also identified below, by checking the box, any foreign application for patent or inventor's certificate or any PCT international application having a filing date before that of the application on which priority is claimed.

Prior Foreign Application Number(s)	Country	Foreign Filing Date (dd,mm,yyyy)	Priority Not Claimed	Certified Copy Attached	
				Yes	No
			<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
			<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

☐ Additional foreign application numbers are listed on a supplemental priority sheet attached hereto.

2) PROVISIONAL PRIORITY CLAIM: I hereby claim the benefit under Title 35, United States Code §119(e) of any United States provisional application(s) listed below.

Provisional Application Number(s)	Filing Date (dd,mm,yyyy)
60/088,670	09 June 1998 (09.06.98)
60/089,138	11 June 1998 (11.06.98)

☐ Additional provisional application numbers are listed on a supplemental priority sheet attached hereto.

3) U.S./PCT PRIORITY CLAIM: I hereby claim the benefit under Title 35, United States Code, §120 of any United States application or §365(c) of any PCT international application designating the United States of America, listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States or PCT international application in the manner provided by the first paragraph of Title 35, United States Code, §112, I acknowledge the duty to disclose information which is known to me to be material to patentability as defined in Title 37, Code of Federal Regulations, §1.56 which became available between the filing date of the prior application and the national or PCT international filing date of this application.

U.S. Parent Application Number	PCT Parent Number	Parent Filing Date (dd,mm,yyyy)	Parent Patent Number (if applicable)
	PCT/US99/12929	09 June 1999 (09.06.99)	

☐ Additional U.S. or PCT international application numbers are listed on a supplemental priority sheet attached hereto.

09719277-041301

POWER OF ATTORNEY:

As a named inventor, I hereby appoint the following attorneys and/or agents to prosecute this application and transact all business in the Patent and Trademark Office connected therewith.

James E. Cockfield	Reg. No. <u>19,162</u>	Nicholas P. Triano III	Reg. No. <u>36,397</u>
Thomas V. Smurzynski	Reg. No. <u>24,798</u>	Peter C. Lauro	Reg. No. <u>32,360</u>
Ralph A. Loren	Reg. No. <u>29,325</u>	DeAnn F. Smith	Reg. No. <u>36,683</u>
Giulio A. DeConti, Jr.	Reg. No. <u>31,503</u>	William D. DeVaul	Reg. No. <u>42,483</u>
Ann Lamport Hammitte	Reg. No. <u>34,858</u>	David J. Ridders	Reg. No. <u>43,882</u>
Elizabeth A. Hanley	Reg. No. <u>33,505</u>	Chi Suk Kim	Reg. No. <u>42,728</u>
Amy E. Mandragouras	Reg. No. <u>36,207</u>	Maria Laccotripe Zacharakis	Limited Recognition Under 37 C.F.R. § 10.9(b)
Anthony A. Laurentano	Reg. No. <u>38,220</u>		
Jane E. Remillard	Reg. No. <u>38,872</u>	Debra J. Milasincic	Reg. No. <u>46,931</u>
Jeremiah Lynch	Reg. No. <u>17,425</u>	David R. Burns	Reg. No. <u>46,590</u>
Kevin J. Canning	Reg. No. <u>35,470</u>	Sean D. Detweiler	Reg. No. <u>42,482</u>
Jeanne M. DiGiorgio	Reg. No. <u>41,710</u>	Peter S. Stecher	Reg. No. <u>47,259</u>
Megan E. Williams	Reg. No. <u>43,270</u>	Adam M. Goodmann	Reg. No. <u>43,640</u>

Send Correspondence to:

Giulio A. DeConti, Jr., Lahive & Cockfield, LLP, 28 State Street, Boston, Massachusetts 02109, United States of America

Direct Telephone Calls to: (name and telephone number)

Giulio A. DeConti, Jr., (617) 227-7400

Wherefore I petition that letters patent be granted to me for the invention or discovery described and claimed in the attached specification and claims, and hereby subscribe my name to said specification and claims and to the foregoing declaration, power of attorney, and this petition.

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Full name of sole or first inventor	<u>Andrea D. BRANCH</u>
Inventor's signature	Date <i>Andrea D. Branch</i> <i>April 10, 2001</i>
Residence	<u>923 5th Avenue, Apartment 6A, New York, New York 10021, United States of America</u>
Citizenship	<u>United States of America</u>
Post Office Address (if different)	<u>same as above</u>

NY

2-00

Full name of sole or first inventor Jose L. WALEWSKI	
Inventor's signature <i>Jose L Walewski</i>	Date <i>April 10, 2001</i>
Residence 95 Tuckahoe Avenue, Eastchester, New York 10709, United States of America NY	
Citizenship United States of America	
Post Office Address (if different) same as above	

3-00

Full name of sole or first inventor Dechard D. STUMP	
Inventor's signature <i>Dechard D Stump</i>	Date <i>APRIL 11, 2001</i>
Residence 529 East 83rd Street, Apartment 5R, New York, New York 10028, United States of America NY	
Citizenship United States of America	
Post Office Address (if different) same as above	

0971937 4267260